

For Research Use Only

Cystatin B Polyclonal antibody

Catalog Number: 10823-1-AP

1 Publications



Basic Information

Catalog Number:

10823-1-AP

Size:

150ul , Concentration: 1000 µg/ml by Nanodrop and 560 µg/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1262

GenBank Accession Number:

BC003370

GeneID (NCBI):

1476

Full Name:

cystatin B (stefin B)

Calculated MW:

11 kDa

Observed MW:

11 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:4000

IP 0.5-4.0 µg for 1.0-3.0 mg of total

protein lysate

IF 1:200-1:800

Applications

Tested Applications:

FC, IF, IP, WB, ELISA

Cited Applications:

IF

Species Specificity:

human

Cited Species:

human

Positive Controls:

WB : human saliva tissue, MCF-7 cells, HepG2 cells, human saliva

IP : MCF-7 cells,

IF : MCF-7 cells, HepG2 cells

Background Information

Cystatin B (CSTB), a member of the cystatin superfamily protein, is a stefin that functions as an intracellular thiol protease inhibitor and has been thought to play a role in protecting against the proteases leaking from lysosomes. CSTB plays various functions in a variety of diseases, including epithelial ovarian cancer, colon cancer, and myoclonus epilepsy. Evidence indicates that mutations in CSTB are responsible for the primary defects in patients with progressive myoclonic epilepsy (EPM1). One type of mutation responsible for EPM1 is the expansion in the promoter region of this gene of a CCCCCCCCCG repeat from 2-3 copies to 30-78 copies.

Notable Publications

Author	Pubmed ID	Journal	Application
Emna Ouni	35341935	Matrix Biol	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human saliva were subjected to SDS PAGE followed by western blot with 10823-1-AP (Cystatin B antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using Cystatin B antibody (10823-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Cystatin B antibody (10823-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).



IP Result of anti-Cystatin B (IP:10823-1-AP, 4ug; Detection:10823-1-AP 1:4000) with MCF-7 cells lysate 1320ug.



1X10⁶ HepG2 cells were stained with 0.2ug Cystatin B antibody (10823-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.