For Research Use Only

HNRNPAO Polyclonal antibody

Catalog Number: 10848-1-AP

1 Publications



Basic Information

Catalog Number:

10848-1-AP BC019271

150ul, Concentration: 300 µg/ml by 10949 Nanodrop and 233 µg/ml by Bradford Full Name:

method using BSA as the standard;

Rabbit Isotype: IgG Observed MW: Immunogen Catalog Number:

AG1312

Size:

Applications

Tested Applications: IF, IP, WB, ELISA **Cited Applications:**

IF. WB

mouse

Species Specificity: human, mouse **Cited Species:**

GenBank Accession Number:

GeneID (NCBI):

heterogeneous nuclear ribonucleoprotein A0

Calculated MW:

31 kDa

Positive Controls:

WB: HeLa cells, fetal human brain tissue, mouse brain

Purification Method:

WB 1:1000-1:6000

for WB

IF 1:50-1:500

Antigen affinity purification

IP 0.5-4.0 ug for IP and 1:500-1:1000

Recommended Dilutions:

tissue, Jurkat cells, NIH/3T3 cells

IP: mouse brain tissue, IF: HeLa cells,

Background Information

HNRNPAO belongs to the A/B subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). Two alternatively spliced transcript variants encoding different isoforms have been described for this gene.

Notable Publications

Author Pubmed ID Journal Application WB,IF Xinxin Zuo 33398173 Nat Struct Mol Biol

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



mouse brain tissue were subjected to SDS PAGE followed by western blot with 10848-1-AP (HNRNPAO antibody) at dilution of 1:500.



Various lysates were subjected to SDS PAGE followed by western blot with 10848-1-AP (HNRNPAO antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (10% Formaldehyde) fixed HeLa cells using 10848-1-AP (HNRNPAO antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



HeLa cells were subjected to SDS PAGE followed by western blot with 10848-1-AP (HNRNPAO Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.