

For Research Use Only

# BLNK Polyclonal antibody

Catalog Number: 10855-1-AP **2 Publications**



## Basic Information

<b>Catalog Number:</b> 10855-1-AP	<b>GenBank Accession Number:</b> BC018906	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 900 µg/ml by Nanodrop and 533 µg/ml by Bradford method using BSA as the standard;	<b>GeneID (NCBI):</b> 29760	<b>Recommended Dilutions:</b> WB 1:500-1:2000 IHC 1:50-1:500 IF 1:50-1:500
<b>Source:</b> Rabbit	<b>Full Name:</b> B-cell linker	
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 50 kDa	
<b>Immunogen Catalog Number:</b> AG1290	<b>Observed MW:</b> 68 kDa, 70 kDa	

## Applications

<b>Tested Applications:</b> IF, IHC, WB, ELISA	<b>Positive Controls:</b> WB : Raji cells, Ramos cells
<b>Cited Applications:</b> IHC, WB	<b>IHC :</b> human tonsillitis tissue, human pancreas cancer tissue
<b>Species Specificity:</b> human, mouse	<b>IF :</b> NIH/3T3 cells,
<b>Cited Species:</b> rat, mouse	
<b>Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	

## Background Information

### Notable Publications

Author	Pubmed ID	Journal	Application
Yi Cheng	34521030	Cytokine	IHC
Yuqi Xin	37925175	Aging (Albany NY)	WB

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



Raji cells were subjected to SDS PAGE followed by western blot with 10855-1-AP (BLNK Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed NIH/3T3 cells using 10855-1-AP (BLNK antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 10855-1-AP (BLNK Antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 10855-1-AP (BLNK Antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).