### For Research Use Only

# IBA1 Polyclonal antibody

Catalog Number: 10904-1-AP

**Featured Product** 

338 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number:

10904-1-AP BC009474
Size: Genel D (NCBI):

150ul , Concentration: 500 µg/ml by 199

Nanodrop; Full Name:
Source: allograft infl

Source: allograft inflammatory factor 1
Rabbit Calculated MW:

Isotype: 17 kDa

IgG .

Immunogen Catalog Number:

AG1363

**Applications** 

Tested Applications:

FC, IF, IHC, WB, ELISA

Cited Applications:

IF, IHC

Species Specificity:

human, mouse, rat

**Cited Species:** 

human, rat, sheep, mouse, zebrafish, hamster, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: THP-1 cells,

IHC: rat brain tissue, human tonsillitis tissue, human liver tissue, mouse brain tissue, human brain tissue

**Purification Method:** 

WB 1:500-1:2000 IHC 1:400-1:1600

IF 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

IF: mouse brain tissue, rat brain tissue

## **Background Information**

IBA1 is a 143 amino acid cytoplasmic, inflammation response scaffold protein. It is constitutively expressed in monocytes and macrophages and is known to be involved in macrophage activation. It is a marker of activated macrophage. Expression of IBA1 is up-regulated in activated microglia following facial nerve axotomy, ischemia, and several brain diseases, thereby implicating it in the activated phenotypes of microglia.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yi-Na Zhang	36168082	Transl Stroke Res	IF
Anna Piotrowska	31616413	Front Immunol	
R Zajaczkowska	30267794	Pharmacol Biochem Behav	

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

#### **Selected Validation Data**




Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 10904-1-AP (IBA1 antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




WB result of IBA1 antibody (10904-1-AP; 1:1000; room temperature for 1.5 hours) with wild-type and IBA1 knockout THP-1 cells.




Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 10904-1-AP (IBA1 antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using IBA1 antibody (10904-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10^6 THP-1 cells were intracellularly stained with 0.4 ug Anti-Human IBA1 (10904-1-AP) and CoraLite® 488-Conjugated Affini Pure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 10904-1-AP (IBA1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 10904-1-AP (IBA1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).