For Research Use Only

E2F4 Polyclonal antibody

Catalog Number: 10923-1-AP

Featured Product

14 Publications

GenBank Accession Number:



Basic Information

Catalog Number:

10923-1-AP BC033180
Size: GeneID (NCBI):

150ul , Concentration: 600 ug/ml by 1874 Nanodrop and 300 ug/ml by Bradford UNIPROT ID:

method using BSA as the standard; Q16254
Source: Full Name:

Rabbit E2F transcription factor 4, p107/p130-

Isotype: binding

IgG Calculated MW:
Immunogen Catalog Number: 44 kDa

AG1365 Observed MW: 60-70 kDa

Applications

Tested Applications: WB, IP, IHC, ELISA

Cited Applications: WB, IHC, IF, IP, ChIP Species Specificity: human, mouse, rat

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:**

WB: A431 cells, Jurkat cells, HL-60 cells, Raji cells,

Purification Method:

WB 1:500-1:2000

protein lysate

IHC 1:20-1:200

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

HeLa cells, NIH/3T3 cells

IP: mouse brain tissue,

IHC: human lymphoma tissue,

Background Information

Transcription factor E2F4 (E2F4) is a transcription activator that binds DNA cooperatively with DP proteins through the E2 recognition site 5'-TTTC[CG]CGC-3' found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication. The DRTF1/E2F complex functions in the control of cell-cycle progression from G1 to S phase. E2F-4 binds with high affinity to RBL1 and RBL2. In some instances, can also bind RB protein. The observed molecular weight of 60-70 kDa is the phosphorylation of E2F4 protein.

Notable Publications

Author	Pubmed ID	Journal	Application
Yinghui Wang	34601503	Cell Death Dis	WB
Xiwen Zhang	29987913	J Cell Physiol	WB,IHC
Yuting Wang	35023014	Protein Cell	WB,ChIP

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of E2F4 antibody (10923-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-E2F4 transfected HeLa cells.



A431 cells were subjected to SDS PAGE followed by western blot with 10923-1-AP (E2F4 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human lymphoma using 10923-1-AP (E2F4 antibody) at dilution of 1:100 (under 10x lens)



IP result of anti-E2F4 (IP:10923-1-AP, 3ug; Detection:10923-1-AP 1:1000) with mouse brain tissue lysate 3600ug.