For Research Use Only

AMPK Alpha Polyclonal antibody Catalog Number: 10929-2-AP Featured Product 261 Publica

Featured Product

261 Publications


oroteinter Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 10929-2-AP	GenBank Accession Number: BC012622		Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):		Recommended Dilu	
	150ul , Concentration: 500 ug/ml by	5562		WB 1:2000-1:10000	
	Nanodrop;	UNIPROT ID:		IP 0.5-4.0 ug for 1.0-	
	Source:	Q13131		protein lysate	
	Rabbit	Full Name:		IHC 1:50-1:500	
	Isotype:	protein kinase, AMP-activated, alpha		IF/ICC 1:200-1:800	
	lgG	1 catalytic subunit	1 catalytic subunit		
	Immunogen Catalog Number: AG1367	Calculated MW: 63 kDa			
		Observed MW:			
		64 kDa			
Applications	Tested Applications:		Positive Controls:		
	WB, IHC, IF/ICC, IP, ELISA	WB: HeLa cel		ls, Jurkat cells, NIH/3T3 cells	
	Cited Applications:	IP : HeLa cells		v	
	WB, IHC, IF, IP		IHC : human p	rostate cancer tissue	·,
	Species Specificity: human, mouse, rat		IF/ICC : HepG2 cells,		
	Cited Species:		in nee incpu		
	human, mouse, rat, pig, rabbit, zebrafish, goat, fish,				
	carp				
	•	non rotrioval			
	carp Note-IHC: suggested antion with TE buffer pH 9.0; (*) A antigen retrieval may be p with citrate buffer pH 6.0	gen retrieval Alternatively, performed			
Background Information	Note-IHC: suggested antic with TE buffer pH 9.0; (*) A antigen retrieval may be p	erformed ated protein kinase (Al switching off ATP-con se. It is a mammalian l rreonine protein kinaso	suming biosynth homologue of su e family. It has 2	etic pathways. PRKA crose non-fermenting isoforms with molec	A1 is also named g protein kinase cular mass of 63-
	Note-IHC: suggested antio with TE buffer pH 9.0; (*) A antigen retrieval may be p with citrate buffer pH 6.0 The mammalian 5-prime-AMP-active stresses that cause ATP depletion by AMPK1, ACACA kinase, HMGCR kinase (SNF-1), which belongs to a serine/th kDa produced by alternative splicing	erformed ated protein kinase (Al switching off ATP-con se. It is a mammalian l rreonine protein kinaso	suming biosynth homologue of su e family. It has 2 mize both AMPK	etic pathways. PRKA crose non-fermenting isoforms with molec	A1 is also named g protein kinase cular mass of 63-1
	Note-IHC: suggested antion with TE buffer pH 9.0; (*) A antigen retrieval may be p with citrate buffer pH 6.0 The mammalian 5-prime-AMP-active stresses that cause ATP depletion by AMPK1, ACACA kinase, HMGCR kinas (SNF-1), which belongs to a serine/th kDa produced by alternative splicing Author Put	ated protein kinase (Ai switching off ATP-con se. It is a mammalian l preonine protein kinase (20929-2-AP can recog med ID Journ	suming biosynth homologue of su e family. It has 2 gnize both AMPK	etic pathways. PRKA crose non-fermenting isoforms with molec	A1 is also named g protein kinase cular mass of 63-1 lpha 2.
Background Information	Note-IHC: suggested antiger with TE buffer pH 9.0; (*) A antigen retrieval may be p with citrate buffer pH 6.0 The mammalian 5-prime-AMP-active stresses that cause ATP depletion by AMPK1, ACACA kinase, HMGCR kinase (SNF-1), which belongs to a serine/th kDa produced by alternative splicing Author Put Xiao-Fei Guo Zent 291	erformed ated protein kinase (Al switching off ATP-con se. It is a mammalian l reconine protein kinase (10929-2-AP can recog med ID Journ 113747 Prost	suming biosynth homologue of su e family. It has 2 gnize both AMPK	etic pathways. PRKA. crose non-fermenting isoforms with molec Alpha 1 and AMPK Al	A1 is also named g protein kinase cular mass of 63- lpha 2. Application
	Note-IHC:suggested antigwith TE buffer pH 9.0; (*) Aantigen retrieval may be pwith citrate buffer pH 6.0The mammalian 5-prime-AMP-activestresses that cause ATP depletion byAMPK1, ACACA kinase, HMGCR kinase(SNF-1), which belongs to a serine/thkDa produced by alternative splicingAuthorPutXiao-Fei Guo291Xiaochen Bai345	ated protein kinase (Al switching off ATP-con se. It is a mammalian l areonine protein kinase g.10929-2-AP can recog omed ID Journ 113747 Prosta 543233 Aging	suming biosynth homologue of su e family. It has 2 mize both AMPK nal aglandins Leukor	etic pathways. PRKA. crose non-fermenting isoforms with molec Alpha 1 and AMPK Al Essent Fatty Acids	A1 is also named g protein kinase cular mass of 63- lpha 2. Application WB
	Note-IHC:suggested antiowith TE buffer pH 9.0; (*) Aantigen retrieval may be rwith citrate buffer pH 6.0The mammalian 5-prime-AMP-activestresses that cause ATP depletion byAMPK1, ACACA kinase, HMGCR kinase(SNF-1), which belongs to a serine/thkDa produced by alternative splicingAuthorPutXiao-Fei Guo291Xiaochen Bai345	erformed ated protein kinase (Al switching off ATP-con se. It is a mammalian l reconine protein kinase (10929-2-AP can recog omed ID Journ (13747 Prost (13747 Prost) (13747 Prost) (13747 Prost) (13747 Prost) (13747 Prost) (13748 Envir ter shipment.	suming biosynth homologue of su e family. It has 2 mize both AMPK hal aglandins Leuko g (Albany NY)	etic pathways. PRKA. crose non-fermenting isoforms with molec Alpha 1 and AMPK Al Essent Fatty Acids	A1 is also named g protein kinase cular mass of 63- lpha 2. Application WB WB


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10929-2-AP (AMPK Alpha 1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 10929-2-AP (AMPK Alpha 1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 10929-2-AP (AMPK Alpha 1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-AMPK Alpha (IP:10929-2-AP, 4ug; Detection:10929-2-AP 1:1000) with HeLa cells lysate 1600 ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using AMPK Alpha antibody (10929-2-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).