For Research Use Only

BID Polyclonal antibody

Catalog Number: 10988-1-AP

Featured Product

62 Publications



Basic Information

Catalog Number: 10988-1-AP

GenBank Accession Number:

Purification Method: Antigen affinity purification

Size:

GeneID (NCBI):

Recommended Dilutions:

150ul, Concentration: 700 ug/ml by Nanodrop and 227 ug/ml by Bradford $\,$ UNIPROT ID:

WB 1:1000-1:4000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

method using BSA as the standard;

P55957

BC009197

protein lysate

Source:

AG1445

Full Name:

Rabbit BH3 interacting domain death agonist

Isotype: Immunogen Catalog Number:

Calculated MW: 22 kDa Observed MW:

22 kDa

Applications

Tested Applications:

WB, IP, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human

Cited Species:

human, mouse, rat, pig

Positive Controls:

WB: A431 cells, human brain tissue, Tunicamycin treated HeLa cells, A549 cells, HEK-293 cells, HepG2

cells, Jurkat cells

IP: HeLa cells,

Background Information

BID, also named as p22 BID, can be cleaved into p11 BID, p13 BID and p15 BID. It is pro-apoptotic molecules. The major proteolytic product p15 BID allows the release of cytochrome c. BID-L, BID-EL and BID-ES induce ICE-like proteases and apoptosis. BID-S does not induce apoptosis. BID counters the protective effect of Bcl-2. (PMID:14583606)

Notable Publications

Author	Pubmed ID	Journal	Application
Bin Xu	31637186	Int J Ophthalmol	WB
Tahir Mehmood	29151958	J Cancer	WB
Robert Pedley	33067418	Cell Death Dis	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10988-1-AP (BID antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



WB result of BID antibody (10988-1-AP, 1:8000) with si-Control and si-BID transfected HeLa cells.



Tunicamycin treated HeLa cells were subjected to SDS PAGE followed by western blot with 10988-1-AP (BID Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-BID (IP:10988-1-AP, 4ug; Detection:10988-1-AP 1:500) with HeLa cells lysate 1280 ug.