

For Research Use Only

# SMARCD1 Polyclonal antibody

Catalog Number: 10998-2-AP

Featured Product

9 Publications



## Basic Information

<b>Catalog Number:</b> 10998-2-AP	<b>GenBank Accession Number:</b> BC009368	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 433 µg/ml by Bradford method using BSA as the standard;	<b>GeneID (NCBI):</b> 6602	<b>Recommended Dilutions:</b> WB 1:500-1:1000 IP 0.5-4.0 ug for IP and 1:500-1:1000 for WB
<b>Source:</b> Rabbit	<b>Full Name:</b> SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily d, member 1	<b>IHC 1:20-1:200</b> <b>IF 1:20-1:200</b>
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 58 kDa	
<b>Immunogen Catalog Number:</b> AG1453	<b>Observed MW:</b> 58 kDa	

## Applications

### Tested Applications:

IF, IHC, IP, WB, ELISA

### Cited Applications:

CoIP, IHC, WB

### Species Specificity:

human, mouse, rat

### Cited Species:

human, mouse

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

WB : mouse brain tissue, human brain tissue, Jurkat cells

IP : mouse brain tissue,

IHC : human lymphoma tissue,

IF : SH-SY5Y cells,

## Background Information

SMARCD1 belongs to the nuclear SWI/SNF chromatin remodeling complex, which regulates transcription and interacts with nucleolin in the nucleus. In addition, it's a member of the neural progenitors-specific chromatin remodeling complex (npBAF complex) and the neuron-specific chromatin remodeling complex (nBAF complex). It possesses a SWIB domain which is homologous to and shares a common fold with the p53-binding domain of MDM2, a cell cycle regulator which interacts with NPM1. SMARCD1 has a strong influence on the Vitamin D-mediated transcriptional activity from an enhancer Vitamin D receptor element (VDRE). And it can mediate critical interactions between nuclear receptors and the BRG1/SMARCA4 chromatin-remodeling complex for transactivation. This antibody may have a cross reaction with SMARCD2.

## Notable Publications

Author	Pubmed ID	Journal	Application
Xu Zhu	32459350	Nucleic Acids Res	CoIP
Tongyu Liu	32404872	Nat Commun	WB
Shuo Chen	25819031	Cancer Lett	WB

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




mouse brain tissue were subjected to SDS PAGE followed by western blot with 10998-2-AP (SMARCD1 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human lymphoma using 10998-2-AP (SMARCD1 antibody) at dilution of 1:50 (under 10x lens).



Immunofluorescent analysis of SH-SY5Y cells, using SMARCD1 antibody 10998-2-AP at 1:50 dilution and FITC-labeled donkey anti-rabbit IgG (green). Blue pseudocolor = DAPI (fluorescent DNA dye).



IP Result of anti-SMARCD1 (IP:10998-2-AP, 3ug; Detection:10998-2-AP 1:600) with mouse brain tissue lysate 4000ug.