For Research Use Only

PPAN Polyclonal antibody Catalog Number: 11006-1-AP Featured Product

Featured Product






Basic Information	Catalog Number: 11006-1-AP	GenBank Accession Number: BC009833	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	150ul , Concentration: 650 ug/ml by		WB 1:1000-1:4000	
	Nanodrop and 327 ug/ml by Bradford	UNIPROT ID:	IHC 1:500-1:2000	
	method using BSA as the standard;	Q9NQ55		
	Source:	Full Name:		
	Rabbit	peter pan homolog (Drosophila)		
	Isotype:	Calculated MW:		
	IgG	53 kDa		
	Immunogen Catalog Number:	Observed MW:		
	AG1419	55-58 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IHC, ELISA	WB : He	/B : HeLa cells, mouse heart tissue, rat heart tissue, rat ver tissue	
	Cited Applications:	liverti		
	WB, IP, IF, IHC	B, IP, IF, IHC IHC : human colo		
	Species Specificity:			
	human, mouse, rat			
	Cited Species: human, mouse, xenopus			
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
	PPAN (or SSF, SSF1 or SSF2) gene is widely expressed, and encodes peter pan homolog protein which may be localized in nucleus. Two isoforms are produced through alternative splicing. So far the characterization of this protein is largely unknown, multiple phosphorylation sites has been revealed based on high-throughput phosphorylation analysis. This antibody is raised against the C-terminal of huaman PPAN and recognize bands around 55 kDa.			
Background Information	localized in nucleus. Two isoforms ar protein is largely unknown, multiple phosphorylation analysis. This antibu	e produced through alternative phosphorylation sites has been	splicing. So far the characterization of this revealed based on high-throughput	
Background Information	localized in nucleus. Two isoforms ar protein is largely unknown, multiple phosphorylation analysis. This antibu around 55 kDa.	e produced through alternative phosphorylation sites has been	splicing. So far the characterization of this revealed based on high-throughput	
	localized in nucleus. Two isoforms ar protein is largely unknown, multiple phosphorylation analysis. This antibu around 55 kDa.	e produced through alternative phosphorylation sites has been ody is raised against the C-term	splicing. So far the characterization of this revealed based on high-throughput inal of huaman PPAN and recognize bands	
	localized in nucleus. Two isoforms ar protein is largely unknown, multiple phosphorylation analysis. This antibu around 55 kDa. Author Put Tecza Aleksandra A 217	e produced through alternative phosphorylation sites has been ody is raised against the C-term omed ID Journal	splicing. So far the characterization of this revealed based on high-throughput inal of huaman PPAN and recognize bands Application	
	localized in nucleus. Two isoforms ar protein is largely unknown, multiple phosphorylation analysis. This antibu around 55 kDa. Author Put Tecza Aleksandra A 211 Astrid S Pfister 255	e produced through alternative phosphorylation sites has been ody is raised against the C-term omed ID Journal 770895 Biol Cell	splicing. So far the characterization of this revealed based on high-throughput inal of huaman PPAN and recognize bands Application WB,IHC	
	localized in nucleus. Two isoforms ar protein is largely unknown, multiple phosphorylation analysis. This antibu around 55 kDa. Author Put Tecza Aleksandra A 211 Astrid S Pfister 255	e produced through alternative phosphorylation sites has been ody is raised against the C-term pmed ID Journal 770895 Biol Cell 759387 J Biol Chem 558383 Development eer shipment.	splicing. So far the characterization of this revealed based on high-throughput inal of huaman PPAN and recognize bands Application WB,IHC WB, IF	


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 11006-1-AP (PPAN antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



WB result of PPAN antibody (11006-1-AP; 1:12000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PPAN transfected HEK-293 cells.



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 11006-1-AP (PPAN antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).