

## CDK4 Polyclonal antibody

Catalog Number: 11026-1-AP

Featured Product

339 Publications

## Basic Information

<b>Catalog Number:</b> 11026-1-AP	<b>GenBank Accession Number:</b> BC010153	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 450 µg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 1019	<b>Recommended Dilutions:</b> WB 1:1000-1:8000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500 IF 1:50-1:500
<b>Source:</b> Rabbit	<b>Full Name:</b> cyclin-dependent kinase 4	
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 34 kDa	
<b>Immunogen Catalog Number:</b> AG1505	<b>Observed MW:</b> 34 kDa	

## Applications

<b>Tested Applications:</b> FC, IF, IHC, IP, WB, ELISA	<b>Positive Controls:</b>
<b>Cited Applications:</b> CoIP, IF, IHC, IP, RIP, WB	<b>WB :</b> HeLa cells, rat pancreas tissue, HepG2 cells, MCF-7 cells
<b>Species Specificity:</b> human, rat	<b>IP :</b> HeLa cells,
<b>Cited Species:</b> human, rat, sheep, mouse, zebrafish, pig, canine	<b>IHC :</b> human breast cancer tissue, human lung cancer tissue
<b>Note-IHC:</b> suggested antigen retrieval with <b>TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	<b>IF :</b> MCF-7 cells,

## Background Information

Cyclin-dependent kinase-4 (CDK4) is a protein-serine kinase involved in the cell cycle. It is essential for the G1- to S-phase transition during the cell cycle and its expression is primarily controlled at the transcriptional level (PMID:17253961). CCND1-CDK4 axis is not only critical in glial tumor cells but also in stromal-derived cells in the surrounding tumor microenvironment that are vital to sustain tumor outgrowth (PMID:21844184).

## Notable Publications

Author	Pubmed ID	Journal	Application
Yuanyuan Bai	36213833	J Oncol	WB
Taiwei Wang	36169181	Oncol Rep	WB
Yue Wang	27690088	Nutrients	WB

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
 Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
 T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
 E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
 W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11026-1-AP (CDK4 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 11026-1-AP (CDK4 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




WB result of CDK4 antibody (11026-1-AP; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CDK4 transfected HeLa cells.



IP Result of anti-CDK4 (IP:11026-1-AP, 3ug; Detection:11026-1-AP 1:1000) with HeLa cells lysate 4000ug.



1X10<sup>6</sup> MCF-7 cells were stained with 0.2ug CDK4 antibody (11026-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using CDK4 antibody (11026-1-AP) at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).