## For Research Use Only

## RDH11 Polyclonal antibody

Catalog Number: 11077-1-AP



**Basic Information** 

Catalog Number: GenBank Accession Number:

 11077-1-AP
 BC011727

 Size:
 GeneID (NCBI):

150ul , Concentration: 500 ug/ml by S1109
Nanodrop and 500 ug/ml by Bradford method using BSA as the standard; Q8TC12

Source: Full Name:
Rabbit retinol dehydrogenase 11 (all-

Isotype:trans/9-cis/11-cis)IgGCalculated MW:Immunogen Catalog Number:35 kDaAG1485Observed MW:

35 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

IHC 1:50-1:500 IF/ICC 1:200-1:800

**Applications** 

Tested Applications: IHC, IF/ICC, ELISA

Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

IHC: human prostate cancer tissue,

IF/ICC : HaCaT cells,

## **Background Information**

RDH11(Retinol dehydrogenase 11) is also named as ARSDR1(androgen-regulated short-chain dehydrogenase/reductase 1), PSDR1(prostate short-chain dehydrogenase/reductase 1) and belongs to the short-chain dehydrogenases/reductases (SDR) family. RDH11 is involved in steroid synthesis and/or degradation in normal and neoplastic prostate epithelium, and, as such, it may be a key enzyme involved in maintaining intracellular balance of steroid hormones in these cells(PMID:11245473). The deduced protein contains 318 amino acids, and the transcript contains 2 potential polyadenylation signals. It has 2 isoforms produced by alternative splicing.

Storage

Storage:

Storage Ruffer:


Storage Buffe

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 11077-1-AP (RDH11 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HaCaT cells using RDH11 antibody (11077-1-AP) at dilution of 1:400 and CoraLite®594-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-4).