

For Research Use Only

ACVR1 Polyclonal antibody

Catalog Number: 11108-1-AP



Basic Information

Catalog Number: 11108-1-AP	GenBank Accession Number: BC033867	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 600 µg/ml by Nanodrop;	GeneID (NCBI): 90	Recommended Dilutions: WB 1:500-1:1000
Source: Rabbit	Full Name: activin A receptor, type I	
Isotype: IgG	Calculated MW: 509 aa, 57 kDa	
Immunogen Catalog Number: AG1556	Observed MW: 57 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : mouse kidney tissue,
Species Specificity: human, mouse, rat	

Background Information

ACVR1 (activin receptor type I), also known as ALK2 or ACTRI, is a receptor for activin. It forms a stable complex with type II receptor after ligand binding. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling, and type II receptors are required for binding ligands and for expression of type I receptors. ACVR1 is expressed in many tissues including skeletal muscle and chondrocytes. It functions as a receptor for bone morphogenetic protein (BMP) and induces Indian hedgehog in chondrocytes during skeletal development. Mutations in ACVR1 gene are associated with fibrodysplasia ossificans progressive (PMID: 16642017).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse kidney tissue were subjected to SDS PAGE followed by western blot with 11108-1-AP (ACVR1 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.