

For Research Use Only

TAP1 Polyclonal antibody

Catalog Number: 11114-1-AP

Featured Product

28 Publications



Basic Information

Catalog Number:

11114-1-AP

Size:

150ul, Concentration: 700 ug/ml by Nanodrop and 387 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1619

GenBank Accession Number:

BC014081

GeneID (NCBI):

6890

UNIPROT ID:

Q03518

Full Name:

transporter 1, ATP-binding cassette, sub-family B (MDR/TAP)

Calculated MW:

81 kDa

Observed MW:

70-81 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:6000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:20-1:200

IF/ICC 1:20-1:200

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF, IP, RIP

Species Specificity:

human, mouse

Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HCT 116 cells, HepG2 cells, mouse skeletal muscle tissue, HeLa cells, mouse spleen tissue, SW480 cells

IP: HepG2 cells,

IHC: human pancreas cancer tissue,

IF/ICC: HepG2 cells,

Background Information

TAP1, also known as ABCB2, PSF1 or RING4, is a member of the ATP-binding cassette (ABC) family of transmembrane transporters and is an essential component of the major histocompatibility complex (MHC) class I antigen-presenting pathway. TAP is involved in the transport of antigens from the cytoplasm to the endoplasmic reticulum for association with MHC class I molecules. It also acts as a molecular scaffold for the final stage of MHC class I folding. Defects in TAP1 are a cause of bare lymphocyte syndrome type 1 (BLS1). Western blot analysis using this antibody detected a major band around 70-80 kDa in HeLa cells.

Notable Publications

Author	Pubmed ID	Journal	Application
Nima Attaran	36276482	Oncol Lett	IHC
Yan Li	36452477	Clin Transl Immunology	WB
Zhen-Da Wang	36419887	Front Oncol	WB,IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11114-1-AP (TAP1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




WB result of TAP1 antibody (11114-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-TAP1 transfected HeLa cells.




Immunofluorescent analysis of HepG2 cells, using TAP1 antibody 11114-1-AP at 1:50 dilution and Rhodamine-labeled goat anti-rabbit IgG (red). Blue pseudocolor = DAPI (fluorescent DNA dye).




Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 11114-1-AP (TAP1 Antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 11114-1-AP (TAP1 Antibody) at dilution of 1:50 (under 40x lens).



Various lysates were subjected to SDS PAGE followed by western blot with 11114-1-AP (TAP1 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



IP result of anti-TAP1 (IP:11114-1-AP, 4ug; Detection:11114-1-AP 1:3000) with HepG2 cells lysate 1320 ug.