

For Research Use Only

EIF3E Polyclonal antibody

Catalog Number: 11352-1-AP



Basic Information

Catalog Number: 11352-1-AP	GenBank Accession Number: BC000734	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 300 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 3646	Recommended Dilutions: WB 1:500-1:2400 IP 0.5-4.0 ug for IP and 1:500-1:1000 for WB
Source: Rabbit	Full Name: eukaryotic translation initiation factor 3, subunit E	IHC 1:50-1:500 IF 1:10-1:100
Isotype: IgG	Calculated MW: 48 kDa	
Immunogen Catalog Number: AG1856	Observed MW: 48-52 kDa	

Applications

Tested Applications:

IF, IHC, IP, WB, ELISA

Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells

IP : HepG2 cells,

IHC : human breast cancer tissue,

IF : HepG2 cells,

Background Information

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 11352-1-AP (EIF3E antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0)).



HeLa cells were subjected to SDS PAGE followed by western blot with 11352-1-AP (EIF3E antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



IP result of anti-EIF3E (IP:11352-1-AP, 4ug; Detection:11352-1-AP 1:500) with HepG2 cells lysate 880 ug.



Immunofluorescent analysis of HepG2 cells using 11352-1-AP (EIF3E antibody) at dilution of 1:25 and Rhodamine-Goat anti-Rabbit IgG.