For Research Use Only

PSMD12 Polyclonal antibody

Catalog Number: 11412-1-AP

Featured Product

2 Publications



Basic Information

Catalog Number: GenBank Accession Number:

 11412-1-AP
 BC019062

 Size:
 GeneID (NCBI):

150ul , Concentration: 700 µg/ml by 5718

Nanodrop; Full Nam

Source: proteasome (prosome, macropain)
Rabbit 26S subunit, non-ATPase, 12

Isotype: Calculated MW:
IgG 456 aa, 53 kDa
Immunogen Catalog Number: Observed MW:
AG1957 50-55 kDa

Applications

Tested Applications:

IHC, WB, ELISA
Cited Applications:

IHC, WB

Species Specificity: human, mouse, rat Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: MCF-7 cells, mouse lung tissue, rat lung tissue

Purification Method:

WB 1:500-1:3000 IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

IHC: human gliomas tissue,

Background Information

Proteasome 26S subunit, non-ATPase 12 (PSMD12; also known as RPN5) was first discovered in fission yeast. It is a non-ATPase subunit of the 19S regulator of the 26S proteasome complex, which is involved in regulating the cell cycle, repairing DNA damage, and regulating apoptosis by mediating the internal protein balance through the removal of misfolded proteins or damaged intracellular substructures. In addition, PSMD12 contributes to tumor cell growth and apoptosis in breast cancer. (PMID: 33987398)

Notable Publications

Author	Pubmed ID	Journal	Application
Xinna Du	32222279	Biochem Biophys Res Commun	WB
Zhongyong Wang	33987398	Ann Transl Med	WB,IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 11412-1-AP (PSMD12 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 11412-1-AP (PSMD12 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).