#### For Research Use Only

# COX7A1 Polyclonal antibody

Catalog Number: 11413-1-AP

4 Publications



**Basic Information** 

Catalog Number: 11413-1-AP

BC002757 GeneID (NCBI):

**Purification Method:** Antigen affinity purification Recommended Dilutions: WB 1:500-1:2000

150ul, Concentration: 400 µg/ml by 1346 Nanodrop and 227 µg/ml by Bradford Full Name:

IHC 1:20-1:200

method using BSA as the standard;

cytochrome c oxidase subunit VIIa polypeptide 1 (muscle)

GenBank Accession Number:

Rabbit Calculated MW: 79 aa, 9 kDa Isotype: IgG Observed MW: Immunogen Catalog Number: 7-9 kDa

AG1981

**Applications** 

**Tested Applications:** IHC, WB,ELISA

**Cited Applications:** IF, IHC, WB

Species Specificity: human, mouse **Cited Species:** human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

**Positive Controls:** 

WB: mouse heart tissue, human brain tissue, human

kidney tissue

IHC: human gliomas tissue,

## **Background Information**

Cytochrome c oxidase(COX) catalyzes the transfer of electrons from reduced cytochrome c to molecular oxygen. Subunit VIIa of mammalian COX exists in at least 2 isoforms, liver and muscle. COX7A1(Cytochrome c oxidase subunit 7A1, mitochondrial) is also named as COX7AH, cytochrome c oxidase subunit VIIa-H, cytochrome c oxidase subunit VIIa-M and is the muscle isoform of COX subunit VIIa. Northern-blot analysis of primate tissues demonstrated that COXVIIa-M mRNA is present only in muscle tissues(PMID:1327965).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Ben Shabat Moti	34179944	Cereb Cortex	IHC
Veronika Boczonadi	25666558	Int J Biochem Cell Biol	WB, IF
Misako Tanaka	38097064	Biochim Biophys Acta Mol Cell Res	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


\*\*\* 20ul sizes contain 0.1% BSA

Aliquoting is unnecessary for -20°C storage

### Selected Validation Data

 $40 kd \rightarrow$   $30 kd \rightarrow$   $20 kd \rightarrow$   $15 kd \rightarrow$   $10 kd \rightarrow$   $5 kd \rightarrow$ 

mouse heart tissue were subjected to SDS PAGE followed by western blot with 11413-1-AP (COX7A1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human gliomas using 11413-1-AP (COX7A1 antibody) at dilution of 1:100 (under 10x lens).