For Research Use Only

BUB1B Polyclonal antibody

Catalog Number: 11504-2-AP

Featured Product

18 Publications




GenBank Accession Number: **Purification Method:** Catalog Number: **Basic Information** 11504-2-AP BC018739 Antigen affinity purification GenelD (NCBI): Recommended Dilutions: Size: 150ul , Concentration: 300 ug/ml by 701 WB 1:500-1:1000 IHC 1:250-1:1000 Nanodrop: UNIPROT ID: IF/ICC 1:50-1:500 Source 060566 Rabbit Full Name: Isotype budding uninhibited by benzimidazoles 1 homolog beta lgG (yeast) Immunogen Catalog Number: Calculated MW: AG2053 1050 aa, 120 kDa Observed MW: 120-130 kDa **Applications Tested Applications: Positive Controls:** WB, IHC, IF/ICC, ELISA WB: mouse testis tissue, K-562 cells, HeLa cells, **Cited Applications:** human testis tissue WB. IHC. IF IHC : human cervical cancer tissue, human colon **Species Specificity:** cancer tissue human, mouse IF/ICC : SKOV-3 cells. **Cited Species:** human, mouse Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Background Information** BUB1B(Mitotic checkpoint serine/threonine-protein kinase BUB1 beta), also named as BUBR1, MAD3Land SSK1, belongs to the BUB1 subfamily. Human BUB1B is a 120 kDa protein that plays a central role in the spindle assembly checkpoint (SAC), the evolutionary conserved and self-regulatory system of higher organisms that monitors and repairs defects in chromosome segregation in mitotic cells(PMID:22201036). In addition to phosphorylation, BUB1B is also subjected to posttranslational modifications including acetylation(PMID:22167194). Defects in BUB1B are the cause of premature chromatid separation trait (PCS) and mosaic variegated aneuploidy syndrome type 1 (MVA1). Notable Publications Author Pubmed ID Journal Application Front Pharmacol WB Xie Mengyan 36176446 Zhi-Chuan Zhu 29520102 Oncogene WB Nucleic Acids Res Fengjie Yuan 31291454 WB Storage Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage *** 20ul sizes contain 0.1% BSA For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA)


W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 11504-2-AP (BubR1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




WB result of BubR1 antibody (11504-2-AP; 1:300; incubated at room temperature for 1.5 hours) with sh-Control and sh-BubR1 transfected HeLa cells.



mouse testis tissue were subjected to SDS PAGE followed by western blot with 11504-2-AP (BubR1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed SKOV-3 cells using BubR1 antibody (11504-2-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 11504-2-AP (BubR1 antibody at dilution of 1:200 (under 40x lens).