## For Research Use Only

## DPP9 Polyclonal antibody

Catalog Number: 11568-1-AP



**Basic Information** 

Catalog Number:

11568-1-AP BC037948

GenBank Accession Number:

GeneID (NCBI):

150ul , Concentration: 267 µg/ml by 91039 Bradford method using BSA as the **UNIPROT ID:** 

standard; Q86T12

Source: Full Name:

Rabbit dipeptidyl-peptidase 9

Isotype: Calculated MW: 863 aa, 98 kDa Immunogen Catalog Number: Observed MW:

AG2145 98 kDa

**Applications Tested Applications:** 

WB, IF, IHC, ELISA Species Specificity:

Note-IHC: suggested antigen retrieval with

TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

human, mouse

Positive Controls:

WB: HepG2 cells, A2780 cells, MCF-7 cells, mouse

**Purification Method:** 

WB 1:500-1:1000 IHC 1:50-1:500

IF/ICC 1:10-1:100

Antigen affinity purification

Recommended Dilutions:

heart tissue, PC-3 cells

IHC: human ovary tumor tissue,

IF/ICC: HepG2 cells, MCF-7 cells, PC-3 cells

**Background Information** 

Dipeptidyl peptidase 9(DPP9), is a member of the S9B family in clan SC of the serine proteases. DPP9 has 3 isoforms with MW of 95, 98 and 101 kDa. DPP9 is involved in the regulation of the activity of their substrates and have been linked to a variety of diseases including type 2 diabetes, obesity and cancer.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**




HepG2 cells were subjected to SDS PAGE followed by western blot with 11568-1-AP (DPP9 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 11568-1-AP (DPP9 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 11568-1-AP (DPP9 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of HepG2 cells using 11568-1-AP (DPP9 antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).