

For Research Use Only

Podoplanin Polyclonal antibody

Catalog Number: 11629-1-AP **14 Publications**



Basic Information

Catalog Number: 11629-1-AP	GenBank Accession Number: BC022812	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 600 µg/ml by Nanodrop;	GeneID (NCBI): 10630	Recommended Dilutions: WB 1:500-1:2000 IHC 1:400-1:1600
Source: Rabbit	UNIPROT ID: Q86YL7	
Isotype: IgG	Full Name: podoplanin	
Immunogen Catalog Number: AG2201	Calculated MW: 17 kDa, 25 kDa	
	Observed MW: 43 kDa	

Applications

Tested Applications: FC, IHC, WB, ELISA	Positive Controls: WB : HEK-293 cells, human placenta tissue IHC : human tonsillitis tissue, human appendicitis tissue
Cited Applications: IF, IHC, WB	
Species Specificity: human	
Cited Species: human, rat	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Podoplanin was identified as a 43 kDa glycoprotein found in the cell membranes of glomerular epithelial cells (podocyte) (PMID: 9327748). It is a lymphatic marker because the expression of podoplanin has been detected in lymphatic but not blood vascular endothelium, and is useful as the marker of tumor-associated Lymphangiogenesis. Podoplanin has a function in developing testis, most likely at the level of cell-cell interactions among pre-meiotic germ cells and immature Sertoli cells. It may be involved in cell migration and/or actin cytoskeleton organization. When expressed in keratinocytes, PDPN induces changes in cell morphology with transfected cells showing an elongated shape, numerous membrane protrusions, major reorganization of the actin cytoskeleton, increased motility and decreased cell adhesion. It is required for normal lung cell proliferation and alveolus formation at birth. PDPN induces platelet aggregation. It does not have any effect on folic acid or amino acid transport and does not function as a water channel or as a regulator of aquaporin-type water channels. The antibody is specific to Podoplanin.

Notable Publications

Author	Pubmed ID	Journal	Application
Yao Ding	34492084	PLoS Pathog	WB,IHC
Wenkun Fu	33995679	Theranostics	IHC
Xiaomin Zhang	24527067	Oncol Lett	IHC

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HEK-293 cells were subjected to SDS PAGE followed by western blot with 11629-1-AP (Podoplanin antibody at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 11629-1-AP (Podoplanin antibody) at dilution of 1:800 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 11629-1-AP (Podoplanin antibody) at dilution of 1:800 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10⁶ HUVEC cells were stained with 0.2ug Podoplanin antibody (11629-1-AP, red) and control antibody (blue). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1500. Cells were fixed with 4% PFA and permeabilized with 0.1% Triton X-100.