For Research Use Only

PKIA Polyclonal antibody

Catalog Number:11743-1-AP



Purification Method:

IHC 1:50-1:500

IF/ICC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number:

11743-1-AP BC022265 GeneID (NCBI): Size:

150ul, Concentration: 650 ug/ml by Nanodrop and 347 ug/ml by Bradford $\,$ UNIPROT ID: method using BSA as the standard; P61925

Source: Full Name: Rabbit

protein kinase (cAMP-dependent, Isotype: catalytic) inhibitor alpha

Calculated MW: Immunogen Catalog Number: 76 aa. 8 kDa

AG2338

Applications Tested Applications:

IHC, IF/ICC, ELISA IHC: mouse embryo tissue, mouse brain tissue

Species Specificity: IF/ICC: HeLa cells, human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

cAMP-dependent protein kinase inhibitor alpha (PKIA) is also named as PRKACN1, and belongs to the PKI family. Among them, PKIA, a member of protein kinase A (PKA) inhibitor family, was found to be most significantly and highly expressed in susceptible animals (PMID:24910983). Decreased PKIA signaling would directly impact PKA activation, potentially inducing DRP1 phosphorylation and increasing ER Ca2+ stores (PMID:32152556). miR-129-5p activated PKA to regulate the phosphorylation of beta-catenin and cAMP-response element binding protein (CREB) by inhibiting PKIA (PMID:36222334).

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data




Immunohistochemical analysis of paraffinembedded mouse embryo tissue slide using 11743-1-AP (PKIA antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse embryo tissue slide using 11743-1-AP (PKIA antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 11743-1-AP (PKIA antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 11743-1-AP (PKIA antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using PKIA antibody (11743-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).