

For Research Use Only

FAPA Polyclonal antibody

Catalog Number: 11779-1-AP **1 Publications**



Basic Information

Catalog Number: 11779-1-AP	GenBank Accession Number: BC026250	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 300 µg/ml by Nanodrop;	GeneID (NCBI): 2191	Recommended Dilutions: WB 1:500-1:1000
Source: Rabbit	Full Name: fibroblast activation protein, alpha	
Isotype: IgG	Calculated MW: 760 aa, 88 kDa	
Immunogen Catalog Number: AG2439	Observed MW: 87-97 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : U-87 MG cells, U2OS cells
Cited Applications: IF	
Species Specificity: human	
Cited Species: mouse	

Background Information

Fibroblast activation protein-alpha (FAP-alpha) is a cell surface glycoprotein serine protease, which can participate in extracellular matrix degradation and is involved in many cellular processes including tissue remodeling, fibrosis, wound healing, inflammation and tumor growth (PMID: 14751930; 16223769). FAP-alpha is a single transmembrane protein that locates the cytoplasm and plasma membrane. Immunohistochemical studies have shown that FAP-alpha is transiently expressed in certain normal fetal mesenchymal tissues but that normal adult tissues as well as malignant epithelial, neural, and hematopoietic cells are generally FAP-alpha-negative. However, most of the common types of epithelial cancers contain abundant FAP-alpha-reactive stromal fibroblasts (PMID: 7911242).

Notable Publications

Author	Pubmed ID	Journal	Application
Jing Kong	31796058	Mol Cancer	IF

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 11779-1-AP (FAPA antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.