### For Research Use Only

# CRNN Polyclonal antibody

Catalog Number: 11799-1-AP

**Featured Product** 

11 Publications



**Basic Information** 

Catalog Number: 11799-1-AP

GenBank Accession Number:

**Purification Method:** Antigen affinity purification

Size:

BC030807 GeneID (NCBI):

Recommended Dilutions:

150ul, Concentration: 300 µg/ml by

cornulin

49860

WB 1:200-1:1000 IHC 1:20-1:200

Nanodrop and 213 µg/ml by Bradford Full Name: method using BSA as the standard;

Calculated MW: Rabbit 495 aa, 54 kDa Isotype: Observed MW: IgG 70 kDa, 58 kDa

Immunogen Catalog Number:

AG2374

**Applications** 

**Tested Applications:** 

IHC, WB, ELISA

Cited Applications:

IF, IHC, WB

Species Specificity:

human, mouse

**Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: mouse brain tissue, HepG2 cells

IHC: human oesophagus tissue, human cervical cancer

tissue, human skin tissue

## **Background Information**

Cornulin, also known as C1orf10, is a member of the "fused gene" family of proteins, which contain N-terminus EFhand domains and multiple tandem peptide repeats. It may play a role in the mucosal/epithelial immune response and epidermal differentiation. Specific to squamous epithelia cells, cornulin is expressed in esophagus, primary keratinocytes, scalp skin, foreskin, cervix, larynx, and fetal bladder. It may be a marker of late epidermal differentiation, and is down-regulated in eczema.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Changji Li	30009832	J Invest Dermatol	WB,IHC
Hsu Po-Kuei PK	24263008	J Thorac Cardiovasc Surg	IHC
Saira Saleem	33889206	Ecancermedicalscience	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


#### **Selected Validation Data**



mouse brain tissue were subjected to SDS PAGE followed by western blot with 11799-1-AP (CRNN antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human oesophagus tissue slide using 11799-1-AP (CRNN Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieved with Citric acid buffer, pH6.0.



Immunohistochemical analysis of paraffinembedded human oesophagus tissue slide using 11799-1-AP (CRNN Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieved with Citric acid buffer, pH6.0.