

For Research Use Only

SDC4 Polyclonal antibody

Catalog Number: 11820-1-AP

Featured Product

5 Publications



Basic Information

Catalog Number: 11820-1-AP	GenBank Accession Number: BC030805	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 600 µg/ml by Nanodrop;	GeneID (NCBI): 6385	Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500
Source: Rabbit	Full Name: syndecan 4	
Isotype: IgG	Calculated MW: 22 kDa	
Immunogen Catalog Number: AG2388	Observed MW: 22-26 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls: WB : NIH/3T3 cells, IHC : mouse skeletal muscle tissue, human liver cancer tissue
Cited Applications: IF, IHC, WB	
Species Specificity: human, mouse	
Cited Species: human, mouse, rat	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

SDC4, also named as syndecan 4, is a cell surface proteoglycan that bears heparan sulfate, which is involved in normal wound healing. It has been reported that SDC4 may be associated with keratinocyte proliferation and differentiation in the epidermis. SDC4 also plays a key role in the angiogenesis and inflammation. The predicted MW of SDC4 is 22 kDa and 11820-1-AP antibody recognizes a protein around 24-26 kDa and 48-52 kDa (homodimer) which is similar to the papers. (PMID: 20639201, 21949132; 18638267; 30165731)

Notable Publications

Author	Pubmed ID	Journal	Application
Lianhong Yin	29033837	Front Pharmacol	WB, IF
Ting-Ting Liu	36375317	eBioMedicine	WB
Heng Yang	33990545	Cell Death Dis	WB, IF, IHC

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



NIH/3T3 lysates were subjected to SDS PAGE followed by western blot with 11820-1-AP (SDC4 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 11820-1-AP (SDC4 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).