

For Research Use Only

PPP1R2P9 Polyclonal antibody

Catalog Number: 11969-1-AP



Basic Information

Catalog Number: 11969-1-AP	GenBank Accession Number: BC056673	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 800 µg/ml by Nanodrop and 433 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 80316	Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500 IF 1:10-1:100
Source: Rabbit	Full Name: protein phosphatase 1, regulatory (inhibitor) subunit 2 pseudogene 9	
Isotype: IgG	Calculated MW: 202 aa, 23 kDa	
Immunogen Catalog Number: AG2591	Observed MW: 35 kDa	

Applications

Tested Applications:

IF, IHC, WB, ELISA

Species Specificity:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : human testis tissue, human brain tissue

IHC : mouse testis tissue, human brain tissue

IF : HepG2 cells,

Background Information

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 11969-1-AP (PPP1R2P9 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 11969-1-AP (PPP1R2P9 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



human testis tissue were subjected to SDS PAGE followed by western blot with 11969-1-AP (PPP1R2P9 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HepG2 cells, using PPP1R2P9 antibody 11969-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).