

For Research Use Only

Glutamate receptor 2 Polyclonal antibody



Catalog Number: 11994-1-AP

23 Publications

Basic Information

Catalog Number: 11994-1-AP	GenBank Accession Number: BC010574	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 750 µg/ml by Nanodrop and 420 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 2891	Recommended Dilutions: WB 1:500-1:2000 IP 0.5-4.0 ug for IP and 1:500-1:1000 for WB
Source: Rabbit	Full Name: glutamate receptor, ionotropic, AMPA 2	Observed MW: 94-105 kDa
Isotype: IgG	Calculated MW: 883 aa, 99 kDa	
Immunogen Catalog Number: AG2662		

Applications

Tested Applications:

FC, IHC, IP, WB, ELISA

Cited Applications:

IF, IHC, IP, WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse brain tissue, human brain tissue, SKOV-3 cells

IP : mouse brain tissue,

IHC : mouse brain tissue,

Background Information

Glutamate receptors are the predominant excitatory neurotransmitter receptors in the mammalian brain and are activated in a variety of normal neurophysiologic processes. GRIA2 (glutamate receptor 2, GluR-2) belongs to a family of glutamate receptors that are sensitive to alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate (AMPA), and function as ligand-activated cation channels that mediate the fast component of excitatory postsynaptic currents in neurons of the central nervous system. These channels are assembled from 4 related subunits, GRIA1-4. The presence of GRIA2 in the heteromeric AMPA receptors impermeabilizes it to calcium, preventing possible calcium-mediated toxicity.

Notable Publications

Author	Pubmed ID	Journal	Application
YuanYuan Hou	34710402	Exp Neurol	WB
Joanna Gruszczynska-Biegala	27826230	Front Cell Neurosci	WB
Mengmeng Wang	31749684	Front Cell Neurosci	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




mouse brain tissue were subjected to SDS PAGE followed by western blot with 11994-1-AP (Glutamate receptor 2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP Result of anti-Glutamate receptor 2 (IP:11994-1-AP, 4ug; Detection:11994-1-AP 1:500) with mouse brain tissue lysate 4000ug.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 11994-1-AP (Glutamate receptor 2 antibody at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10⁶ SH-SY5Y cells were stained with 0.2ug Glutamate receptor 2 antibody (11994-1-AP, red) and control antibody (blue). Fixed with 4% PFA blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1500.