For Research Use Only

STT3A Polyclonal antibody

Catalog Number: 12034-1-AP

Featured Product





Basic Information

Catalog Number: 12034-1-AP Size: 150ul , Concentration: 500 ug/ml by Nanodrop: Source Rabbit Isotype: lgG Immunogen Catalog Number: AG2698

GenBank Accession Number: BC020965 GenelD (NCBI): 3703 UNIPROT ID: P46977 Full Name: STT3, subunit of the

homolog A (S. cerevisiae)

Calculated MW:

705 aa, 81 kDa Observed MW: 65-100 kDa

oligosaccharyltransferase complex,

Positive Controls:

tissue

WB: HepG2 cells, K-562 cells

IHC : human stomach tissue, human stomach cancer

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500

Applications

Tested Applications: WB, IHC, ELISA **Cited Applications:** WB, IHC, IF, IP, CoIP **Species Specificity:** human **Cited Species:** human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

STT3A, also named as Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit STT3A, is a 705 amino acid prtein, which belongs to the STT3 family. STT3A is expressed at high levels in placenta, liver, muscle and pancreas, and at very low levels in brain, lung and kidney. STT3A is a catalytic subunit of the N-oligosaccharyl transferase (OST) complex which catalyzes the transfer of a high mannose oligosaccharide from a lipid-linked oligosaccharide donor to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent polypeptide chains. N-glycosylation occurs cotranslationally and the complex associates with the Sec61 complex at the channelforming translocon complex that mediates protein translocation across the endoplasmic reticulum (ER). SST3A seems to be involved in complex substrate specificity. STT3A is present in the majority of OST complexes and mediates cotranslational N-glycosylation of most sites on target proteins, while STT3B-containing complexes are required for efficient post-translational glycosylation and mediate glycosylation of sites that have been skipped by STT3A.

Notable Publications

Author	Pubmed ID	Journal	Application
Xinxin Song	32938586	Cancer Discov	WB,IF,IP
Shih-Han Wang	36381324	Am J Cancer Res	
Cecilia Lopez-Sambrooks	27694802	Nat Chem Biol	WB

Storage

Storage Store at -20°C. Stable for one year after shipment.

Storage Buffer PBS with 0.02% sodium azide and 50% glycerol, pH7.3 Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



HepG2 cells were subjected to SDS PAGE followed by western blot with 12034-1-AP (STT3A antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 12034-1-AP (STT3A antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human stomach tissue slide using 12034-1-AP (STT3A antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human stomach tissue slide using 12034-1-AP (STT3A antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).