For Research Use Only

TRAPPC4 Polyclonal antibody

Catalog Number: 12074-1-AP



Basic Information

Catalog Number: GenBank Accession Number:

12074-1-AP BC010866 GeneID (NCBI): Size: 150ul, Concentration: 300 ug/ml by 51399

Nanodrop and 200 ug/ml by Bradford $\ensuremath{\,^{\text{UNIPROT\,ID:}}}$ method using BSA as the standard; Q9Y296

Source: Full Name: Rabbit trafficking protein particle complex 4

Isotype: Calculated MW: 219 aa, 24 kDa Immunogen Catalog Number: Observed MW: 24-27 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:4000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:500-1:2000

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen

retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HL-60 cells, mouse testis tissue, rat testis tissue

IP: HL-60 cells.

IHC: human colon cancer tissue.

IF/ICC: HeLa cells,

Background Information

Transport protein particle (TRAPP, also known as trafficking protein particle), a multimeric guanine nucleotideexchange factor, regulates multiple membrane trafficking pathways (PMID: 20966969). TRAPPC4, also known as synbindin, is a core component of the TRAPP complexes and one of the essential subunits for guanine nucleotide exchange factor activity for Rab1 GTPase (PMID: 31794024). Deficiencies in vesicular transport mediated by TRAPPC4 have been associated with severe syndromic intellectual disability (PMID: 31794024).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12074-1-AP (TRAPPC4 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




IP result of anti-TRAPPC4 (IP:12074-1-AP, 4ug; Detection:12074-1-AP 1:800) with HL-60 cells lysate 1720 ug.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 12074-1-AP (TRAPPC4 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 12074-1-AP (TRAPPC4 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using TRAPPC4 antibody (12074-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).