

For Research Use Only

# UCK1 Polyclonal antibody

Catalog Number: 12271-1-AP **2 Publications**



## Basic Information

<b>Catalog Number:</b> 12271-1-AP	<b>GenBank Accession Number:</b> BC015547	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 700 µg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 83549	<b>Recommended Dilutions:</b> WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IF 1:10-1:100
<b>Source:</b> Rabbit	<b>Full Name:</b> uridine-cytidine kinase 1	
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 277 aa, 31 kDa	
<b>Immunogen Catalog Number:</b> AG2912	<b>Observed MW:</b> 33 kDa	

## Applications

<b>Tested Applications:</b> IF, IP, WB, ELISA	<b>Positive Controls:</b> WB : HEK-293T cells, MCF-7 cells
<b>Cited Applications:</b> WB	<b>IP :</b> MCF-7 cells,
<b>Species Specificity:</b> human, mouse, rat	<b>IF :</b> MCF-7 cells,
<b>Cited Species:</b> human, mouse	

## Background Information

Uridine/cytidine kinase-1 is a pyrimidine ribonucleoside kinase that catalyzes the phosphorylation of uridine and cytidine to form uridine monophosphate (UMP) and cytidine monophosphate (CMP). The deduced 277-amino acid protein has a calculated molecular mass of 31 kDa. UCK1 belongs to the uridine kinase family and UCK have important roles for the phosphorylation of nucleoside analogs that may be important in chemotherapy of cancer.

## Notable Publications

Author	Pubmed ID	Journal	Application
Nao Yoshida-Sakai	34913485	Int J Cancer	WB
Tatsuro Watanabe	38190613	Blood Adv	WB

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 12271-1-AP (UCK1 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



IP Result of anti-UCK1 (IP:12271-1-AP, 4 $\mu$ g; Detection:12271-1-AP 1:1000) with MCF-7 cells lysate 2800 $\mu$ g.



Immunofluorescent analysis of MCF-7 cells using 12271-1-AP (UCK1 antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).