## For Research Use Only

# Sortilin Polyclonal antibody

Catalog Number: 12369-1-AP

**Featured Product** 

11 Publications

BC023542



**Basic Information** 

Catalog Number: GenBank Accession Number:

GeneID (NCBI):

150ul, Concentration: 300 µg/ml by 6272

Nanodrop; Full Name: Source: sortilin 1

Rabbit Calculated MW: Isotype: 831 aa, 92 kDa IgG

Observed MW: Immunogen Catalog Number: 100-110 kDa

AG2997

12369-1-AP

**Tested Applications:** 

FC, IF, IHC, IP, WB, ELISA Cited Applications:

CoIP, IF, IHC, WB Species Specificity: human, mouse, rat

**Cited Species:** 

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

**Purification Method:** 

Antigen affinity purification

Recommended Dilutions: WB 1:2000-1:16000

IP 0.5-4.0 ug for IP and 1:500-1:1000

for WB

IHC 1:500-1:2000 IF 1:200-1:800

**Applications** 

Positive Controls:

WB: mouse brain tissue, human brain tissue, rat brain

tissue

IP: rat brain tissue.

IHC: mouse brain tissue, human brain tissue, rat brain

IF: mouse brain tissue.

## **Background Information**

Sortilin 1 (SORT1), is a trans-Golgi network (TGN) transmembrane protein and is a member of the Vps10p domain receptor family. SORT1 binds a number of unrelated ligands that participate in a wide range of cellular processes. It functions as a sorting receptor in the Golgi compartment and as a clearance receptor on the cell surface.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Hiroyuki Obata	36185767	Exp Ther Med	IF
Keiji Uchiyama	28665987	PLoS Pathog	WB,IF
Noah R Johnson	29203673	Proc Natl Acad Sci U S A	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


### Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12369-1-AP (Sortilin antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 12369-1-AP (Sortilin antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 12369-1-AP (Sortilin antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




IP Result of anti-Sortilin (IP:12369-1-AP, 4ug; Detection:12369-1-AP 1:500) with rat brain tissue lysate 4000ug.



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using Sortilin antibody (12369-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using Sortilin antibody (12369-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10^6 Neuro-2a cells were intracellularly stained with 0.4 ug Anti-Human Sortilin (12369-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgC(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).