

For Research Use Only

SF3B14 Polyclonal antibody

Catalog Number: 12379-1-AP **2 Publications**



Basic Information

Catalog Number: 12379-1-AP	GenBank Accession Number: BC015463	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 600 µg/ml by Nanodrop and 333 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 51639	Recommended Dilutions: WB 1:1000-1:4000
Source: Rabbit	Full Name: splicing factor 3B, 14 kDa subunit	
Isotype: IgG	Calculated MW: 14 kDa	
Immunogen Catalog Number: AG3045	Observed MW: 14 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HeLa cells, HepG2 cells
Cited Applications: WB	
Species Specificity: human, mouse, rat	
Cited Species: human	

Background Information

Pre-mRNA splicing by the human spliceosome requires the sequential recognition by RNA and protein factors of the 5' and 3' splice sites, the branch region and the polypyrimidine tract. This sequence is characterized by the stepwise association of the U1, U2, and U4/U5/U6 snRNP particles [PMID:9476892]. Human p14 (SF3b14), a component of the spliceosomal U2 snRNP, interacts directly with the pre-mRNA branch adenosine within the context of the bulged duplex formed between the pre-mRNA branch region and U2 snRNA. This association occurs early in spliceosome assembly and persists within the fully assembled spliceosome [PMID:21062891].

Notable Publications

Author	Pubmed ID	Journal	Application
Teng Teng	28541300	Nat Commun	WB
Frederike L Harms	37422718	Genet Med	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 12379-1-AP (SF3B14 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.