

PD-ECGF Polyclonal antibody

Catalog Number: 12383-1-AP

6 Publications

Basic Information

Catalog Number:

12383-1-AP

GenBank Accession Number:

BC018160

Purification Method:

Antigen affinity purification

Size:

150ul, Concentration: 300 µg/ml by Nanodrop and 240 µg/ml by Bradford method using BSA as the standard;

GeneID (NCBI):

1890

Recommended Dilutions:

WB 1:500-1:2000

Full Name:

thymidine phosphorylase

IHC 1:50-1:500

Source:

Rabbit

Calculated MW:

482 aa, 50 kDa

Isotype:

IgG

Observed MW:

50 kDa

Immunogen Catalog Number:

AG3052

Applications

Tested Applications:

IHC, WB, ELISA

Positive Controls:

WB: A431 cells, HeLa cells, THP-1 cells

Cited Applications:

FC, IF, IHC, WB

IHC: human breast cancer tissue, human stomach cancer tissue, human ovary tumor tissue

Species Specificity:

human

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with **TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

Background Information

Platelet derived endothelial cell growth factor (PD-ECGF), also known as thymidine phosphorylase (TP), induces migration and angiogenesis in endothelial and tumor cells, and was upregulated in various malignancies compared to that in normal tissues. Interestingly, PD-ECGF has dual effect on tumor development and chemotherapy. It could stimulate cancer cell migration and proliferation. On the other hand, some chemotherapeutic agents (5-fluorouracil, capecitabine, etc.) were converted to their active forms through TP enzymes.

Notable Publications

Author	Pubmed ID	Journal	Application
Ryosuke Tamura	25304388	J Ovarian Res	WB
Yuhan Gu	35314790	Oncogene	WB
Qiong Lai	32123298	Acta Pharmacol Sin	WB, IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 12383-1-AP (PD-ECGF antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 12383-1-AP (PD-ECGF antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 12383-1-AP (PD-ECGF antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 12383-1-AP (PD-ECGF antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 12383-1-AP (PD-ECGF antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).