## For Research Use Only

# PIP4K2A Polyclonal antibody

Catalog Number: 12469-1-AP

3 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 12469-1-AP BC018034

re: GeneID (NCBI):

150ul , Concentration: 600 µg/ml by 5305

Nanodrop; Full Name

Source: phosphatidylinositol-5-phosphate 4-Rabbit kinase, type II, alpha
Isotype: Calculated MW:

IgG 406 aa, 46 kDa Immunogen Catalog Number: Observed MW: AG3152 46 kDa Purification Method: Antigen affinity purification

Recommended Dilutions: WB 1:500-1:2000

IP 0.5-4.0 ug for IP and 1:500-1:1000

for WB IHC 1:50-1:500 IF 1:50-1:500

**Applications** 

**Tested Applications:** 

IF, IHC, IP, WB, ELISA

Cited Applications: IF, IHC, IP, WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: mouse brain tissue, HeLa cells, rat brain tissue

IP: mouse brain tissue,

IHC: human spleen tissue, mouse brain tissue

IF: SH-SY5Y cells, HepG2 cells

# **Background Information**

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Archna Ravi	33984270	Dev Cell	IF,IHC
Mei Zhang	35987479	Int J Biochem Cell Biol	IF,IHC,IP,WB
Joanna Triscott	36724278	Sci Adv	WB,IHC,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


#### Selected Validation Data




mouse brain tissue were subjected to SDS PAGE followed by western blot with 12469-1-AP (PIP4K2A antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




IP Result of anti-PIP4K2A (IP:12469-1-AP, 4ug; Detection:12469-1-AP 1:500) with mouse brain tissue lysate 4400ug.




Immunohistochemical analysis of paraffinembedded human spleen tissue slide using 12469-1-AP (PIP4K2A Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human spleen tissue slide using 12469-1-AP (PIP4K2A Antibody) at dilution of 1:200 (under 40x lens).




Immunofluorescent analysis of (4% PFA) fixed SH-SY5Y cells using 12469-1-AP (PIP4k2A antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




Various lysates were subjected to SDS PAGE followed by western blot with 12469-1-AP (PIP4K2A antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



mouse brain tissue were subjected to SDS PAGE followed by western blot with 12469-1-AP (PIP4K2A antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 12469-1-AP (PIP4K2A antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using PIP4k(2A antibody (12469-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (black).