

For Research Use Only

EDIL3 Polyclonal antibody

Catalog Number: 12580-1-AP

18 Publications



Basic Information

Catalog Number:

12580-1-AP

Size:

150ul, Concentration: 300 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG3274

GenBank Accession Number:

BC030828

GeneID (NCBI):

10085

UNIPROT ID:

O43854

Full Name:

EGF-like repeats and discoidin I-like domains 3

Calculated MW:

52 kDa

Observed MW:

52-55 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IHC 1:50-1:500

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IHC, IF, CoIP, ELISA

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, COLO 320 cells, mouse brain tissue, HepG2 cells, mouse lung tissue, human lung tissue, HeLa cells, K-562 cells

IHC: human pancreas cancer tissue, human lung cancer tissue, human liver cancer tissue

Background Information

EGF-like repeats and discoidin I-like domain-containing protein 3 (EDIL3), also named as DEL1 or integrin-binding protein DEL1, is a 52-kDa extracellular matrix protein produced by endothelial cells in embryos (PMID: 9420328). It is composed of three EGF repeats and two discoidin I-like domains. The second EGF repeat contains an RGD motif. EDIL3 promotes adhesion of endothelial cells through interaction with the alpha-v/beta-3 integrin receptor. It may be involved in vascular remodeling during angiogenesis (PMID: 12840057; 14981004). EDIL3 has also been reported to be an endogenous inhibitor of inflammatory cell recruitment by interfering with the integrin LFA-1-dependent leukocyte-endothelial adhesion (PMID: 19008446).

Notable Publications

Author	Pubmed ID	Journal	Application
Tomoki Maekawa	26374165	Nat Commun	ELISA
Won-Young Kim	31506547	Sci Rep	
Ming-Xuan Feng	25273699	Mol Cancer	WB, IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 12580-1-AP (EDIL3 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 12580-1-AP (EDIL3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 12580-1-AP (EDIL3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).