

For Research Use Only

MICA Polyclonal antibody

Catalog Number: 12619-1-AP **2 Publications**



Basic Information

Catalog Number: 12619-1-AP	GenBank Accession Number: BC016929	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 200 µg/ml by Nanodrop and 173 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 100507436	Recommended Dilutions: WB 1:500-1:1000
Source: Rabbit	Full Name: MHC class I polypeptide-related sequence A	
Isotype: IgG	Calculated MW: 383 aa, 43 kDa	
Immunogen Catalog Number: AG3298	Observed MW: 43 kDa, 50-55 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : K-562 cells, HeLa cells
Cited Applications: WB	
Species Specificity: human	
Cited Species: human	

Background Information

Human MHC class I chain-related genes located within the HLA class I region of chromosome 6 encode MHC class I chain-related A and B (MICA and MICB) (PMID: 11429322). MICA and MICB are stress-inducible surface molecules that are not associated with β 2-microglobulin and do not present peptides (PMID: 9497295). They are expressed in intestinal epithelium and many epithelial tumors (PMID: 10359807). MICA and MICB are ligands for NKG2D which is an activating receptor that is expressed on most natural killer (NK) cells, CD8 $\alpha\beta$ T cells, and $\gamma\delta$ T cells (PMID: 10426993; 11491531).

Notable Publications

Author	Pubmed ID	Journal	Application
Qiulin Wu	38254761	Cancers (Basel)	WB
Xicai Li	36765808	Cancers (Basel)	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



K-562 cells were subjected to SDS PAGE followed by western blot with 12619-1-AP (MICA antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.