For Research Use Only

MICA Polyclonal antibody

Catalog Number:12619-1-AP 2 Publications



Basic Information

Catalog Number: 12619-1-AP

GenBank Accession Number:

Purification Method: Antigen affinity purification

Size:

GeneID (NCBI):

Recommended Dilutions:

150ul, Concentration: 200 ug/ml by

100507436

BC016929

WB 1:500-1:1000

Nanodrop and 173 ug/ml by Bradford $\,$ UNIPROT ID: method using BSA as the standard;

Q29983 Full Name:

Source: Rabbit

MHC class I polypeptide-related

Isotype:

sequence A Calculated MW:

Immunogen Catalog Number:

383 aa, 43 kDa

AG3298

Observed MW:

50-55 kDa

Applications

Tested Applications:

WB, ELISA

Cited Applications:

Species Specificity: human

Cited Species:

human

Positive Controls:

WB: HeLa cells, K-562 cells, SH-SY5Y cells, U266 cells

Background Information

Human MHC class I chain-related genes located within the HLA class I region of chromosome 6 encode MHC class I chain-related A and B (MICA and MICB) (PMID: 11429322). MICA and MICB are stress-inducible surface molecules that are not associated with β 2-microglobulin and do not present peptides (PMID: 9497295). They are expressed in intestinal epithelium and many epithelial tumors (PMID: 10359807). MICA and MICB are ligands for NKG2D which is an activating receptor that is expressed on most natural killer (NK) cells, CD8 $\alpha\beta$ T cells, and $\gamma\delta$ T cells (PMID: 10426993; 11491531).

Notable Publications

Author	Pubmed ID	Journal	Application
Qiulin Wu	38254761	Cancers (Basel)	WB
Xicai Li	36765808	Cancers (Basel)	WB

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 12619-1-AP (MICA antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.