## For Research Use Only

## COG8 Polyclonal antibody, PBS Only

Catalog Number: 12661-1-PBS



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** 

Antigen affinity purification

12661-1-PBS

BC017492

GeneID (NCBI):

100ug, Concentration: 1 mg/ml by Nanodrop:

**UNIPROT ID:** 

Rabbit

Q96MW5 Full Name:

Isotype:

component of oligomeric golgi

IgG

complex 8

Immunogen Catalog Number: AG3341

Calculated MW: 612 aa, 68 kDa

Observed MW: 68 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, Indirect ELISA

Species Specificity:

human, mouse, rat

**Background Information** 

The conserved oligomeric Golgi (COG) complex is an eight subunit (COG1-8) peripheral Golgi membrane heterooligomeric protein complex. The COG complex is organized into lobes A (COG2-4) and B (COG5-7) with COG1 and COG8 bridging these lobes and is thought to play a critical role in vesicle tethering processes (PMID: 17331980). Cog8 localizes exclusively in the Golgi apparatus and is involved in the retrograde Golgi transport of resident proteins responsible for sugar chain (glycan) biosynthesis (PMID: 17220172).

Storage


Storage:

Store at -80°C.


Storage Buffer:

PBS only, pH7.3


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12661-1-AP (COG8 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 12661-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 12661-1-AP (COG8 antibody) at dilution of 1:200 (under 40% lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 12661-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 12661-1-AP (COG8 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 12661-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed A431 cells using COG8 antibody (12661-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red). This data was developed using the same antibody clone with 12661-1-PBS in a different storage buffer formulation.