### For Research Use Only

# PAICS Polyclonal antibody

Catalog Number: 12967-1-AP

**Featured Product** 

11 Publications



**Basic Information** 

Catalog Number: 12967-1-AP

GenBank Accession Number:

BC010273

GeneID (NCBI):

10606

150ul , Concentration: 550 ug/ml by Nanodrop:

**UNIPROT ID:** 

Size:

IgG

AG3621

P22234

Rabbit Isotype:

phosphoribosylamino imidazole

Full Name: carboxylase,

Immunogen Catalog Number:

phosphoribosylamino imidazole

succinocarboxamide synthetase

Calculated MW: 425 aa, 47 kDa

Observed MW:

47 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, ELISA

Cited Applications:

WB, IHC, IF, CoIP

**Species Specificity:** 

human, mouse

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HL-60 cells, mouse heart tissue, HeLa cells, Jurkat cells, K-562 cells, Raji cells, THP-1 cells, mouse brain

**Purification Method:** 

WB 1:1000-1:8000

IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

IHC: human liver tissue, human testis tissue

# **Background Information**

PAICS is also named as ADE2(Multifunctional protein), AIRC, PAIS. It is an important bifunctional enzyme in de novo purine biosynthesis in vertebrate with both 5-aminoimidazole ribonucleotide carboxylase (AIRc) and 4-(Nsuccinylcarboxamide)-5-aminoimidazole ribonucleotide synthetase (SAICARs) activities. It becomes an attractive target for rational anticancer drug design, since rapidly dividing cancer cells rely heavily on the purine de novo pathway for synthesis of adenine and guanine, whereas normal cells favor the salvage pathway (PMID:17224163).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Eunus S Ali	32485148	Mol Cell	WB
Minjun Meng	30097015	Biol Res	WB
Seiya Yamada	32344379	iScience	IF,CoIP

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 12967-1-AP (PAICS antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



1x10^6 HeLa cells were intracellularly stained with 0.25 ug PAICS Polyclonal antibody (12967-1-AP) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 12967-1-AP (PAICS antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 12967-1-AP (PAICS antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).