For Research Use Only

AXL Polyclonal antibody

Catalog Number: 13196-1-AP

Featured Product

34 Publications



Basic Information

Catalog Number:

13196-1-AP

GenBank Accession Number:

BC032229

Size:

GeneID (NCBI):

150ul , Concentration: 500 ug/ml by

UNIPROT ID: P30530

Rabbit

Nanodrop:

Full Name:

Isotype: IgG

AXL receptor tyrosine kinase

Immunogen Catalog Number:

Calculated MW: 894 aa, 98 kDa

AG3898

Observed MW:

140 kDa

Applications

Tested Applications:

WB, IP, ELISA

Cited Applications:

Species Specificity:

human

Cited Species:

human, rat

Antigen affinity purification Recommended Dilutions:

WB 1:1000-1:8000

Purification Method:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

WB, IHC, IF, IP, CoIP

Positive Controls:

WB: DU 145 cells, HeLa cells, HepG2 cells, human liver

tissue, HT-1080 cells

IP: HeLa cells,

Background Information

Axl, also known as Ufo or Ark, is a receptor tyrosine kinase of the TAM (Tyro3, Axl and Mer) family. Axl is a singlepass type I membrane protein containing two fibronectin type-III domains, two Ig-like C2-type domains and one protein kinase domain. Axl mRNA is widely expressed with highest levels in the heart and skeletal muscle. Gas6 has been identified as a ligand for human Axl. Gas6/Axl signaling plays a role in cell survival, cell proliferation, migration and differentiation. It has been reported that 110-120 kDa non-glycosylated AXL protein can be detected(PMID: 31171001). Endogenous Axl is auto-phosphorylated and reacted with a tyrosine-phosphorylated protein of 140 kDa(PMID: 12730092).

Notable Publications

Author	Pubmed ID	Journal	Application
Pei Li	34679238	Chem Biol Drug Des	WB
Shiye Ke	34629012	Pharm Biol	WB
Shan Su	34637814	Virus Res	FC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



HepG2 cells were subjected to SDS PAGE followed by western blot with 13196-1-AP (AXL antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 13196-1-AP (AXL antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



IP result of anti-AXL (IP:13196-1-AP, 4ug; Detection:13196-1-AP 1:3000) with HeLa cells lysate 1320 ug.