For Research Use Only

CA1 Polyclonal antibody

Catalog Number: 13198-2-AP

9 Publications



Basic Information

Catalog Number: GenBank Accession Number: 13198-2-AP BC027890

Size: GeneID (NCBI):

150ul , Concentration: 1000 µg/ml by 759 Nanodrop and 440 µg/ml by Bradford $_{\mbox{Full Name:}}$

method using BSA as the standard;

method using BSA as the standard; carbonic anhydrase I
Source: Calculated MW:
Rabbit 261 aa, 29 kDa

Isotype: Observed MW:
IgG 29 kDa

Immunogen Catalog Number:

AG3932

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:3000

IP 0.5-4.0 ug for IP and 1:500-1:1000

for WB IHC 1:50-1:500

Applications

Tested Applications:

IHC, IP, WB, ELISA

Cited Applications: IF, IHC, IP, WB

Species Specificity: human, mouse, rat

Cited Species: human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: mouse spleen tissue, Jurkat cells, rat spleen

tissue

IP: K-562 cells,

IHC: human breast cancer tissue,

Background Information

CA1(Carbonic anhydrase 1) is also named as CAB and belongs to the alpha-carbonic anhydrase family, which may reside in cytoplasm, in mitochondria, or in secretory granules, or associate with membranes in cell. It forms a large family of genes encoding zinc metalloenzymes of great physiologic importance. Extracellular CA1 mediates hemorrhagic retinal and cerebral vascular permeability through prekallikrein activation(PMID:17259996).

Notable Publications

| Author | Pubmed ID | Journal | Application |
|----------------|-----------|-----------------|-------------|
| Jianming Yang | 36114582 | Microbiome | IF |
| Kazuhiro Tange | 36289244 | Sci Rep | WB,IHC,IP |
| Meili Lu | 30364197 | Front Pharmacol | WB |

Storage

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 13198-2-AP (CA1 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 13198-2-AP (CA1 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 13198-2-AP (CA1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP Result of anti-CA1 (IP:13198-2-AP, 4ug; Detection:13198-2-AP 1:500) with K-562 cells lysate 1200ug.