For Research Use Only

DAO Polyclonal antibody

Catalog Number: 13273-1-AP

4 Publications



Basic Information

Catalog Number: GenBank Accession Number: 13273-1-AP BC029057

ize: GeneID (NCBI):

150ul, Concentration: 300 µg/ml by 1610

Nanodrop; Full I

Source: D-amino-acid oxidase
Rabbit Calculated MW:
Isotype: 347 aa, 39 kDa
IgG Observed MW:

IgG Observed MW:
Immunogen Catalog Number: 39 kDa

AG3973

Positive Controls:

WB: mouse kidney tissue, rat cerebellum tissue, A549

Purification Method:

WB 1:500-1:2000 IHC 1:100-1:400

Antigen affinity purification

Recommended Dilutions:

cells

IHC: human colon tissue,

Applications

Tested Applications:

IHC, WB, ELISA

Cited Applications: IF, IHC, WB

Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

D-amino acid oxidase (DAO), with MW of 39 kDa, is a flavoprotein which uses flavin adenine dinucleotide as its prosthetic group. Its substrates include a wide variety of D-amino acids, but it is inactive on the naturally occurring L-amino acids. DAO may play a role in the pathophysiology of schizophrenia and act as a detoxifying agent which removes D-amino acids that accumulate during aging. DAAO exists in solution as a mixture of monomer-dimer(80 kDa) and even higher molecular aggregates in equilibrium(170 kDa), depending on the enzyme concentration and forms(PMID: 11130179/PMID: 9210639).

Notable Publications

Author	Pubmed ID	Journal	Application
Takahiro Seki	30342995	Mol Cell Neurosci	
Danqing Xin	29751053	Brain Behav Immun	WB
Anastasia V Sudarikova	35081320	Am J Physiol Cell Physiol	IF

Storage

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



mouse kidney tissue were subjected to SDS PAGE followed by western blot with 13273-1-AP (DAO antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 13273-1-AP (DAO Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 13273-1-AP (DAO Antibody) at dilution of 1:200 (under 40x lens).