## For Research Use Only

## CEND1 Polyclonal antibody

Catalog Number: 13280-1-AP



**Basic Information** 

Catalog Number: GenBank Accession Number: 13280-1-AP BC034732

GeneID (NCBI): Size:

150ul , Concentration: 300 µg/ml by 51286 Nanodrop and 133 µg/ml by Bradford

method using BSA as the standard;

differentiation 1 Rabbit Calculated MW: 149 aa, 15 kDa Isotype: IgG Observed MW: Immunogen Catalog Number: 23 kDa

AG4024

Antigen affinity purification Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500

**Purification Method:** 

IF 1:50-1:500

**Applications** 

**Tested Applications:** IF, IHC, WB, ELISA

Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

**Positive Controls:** 

WB: mouse brain tissue, SH-SY5Y cells, human brain

tissue, rat brain tissue

IHC: mouse brain tissue, rat brain tissue

IF: Neuro-2a cells,

## **Background Information**

Cell cycle exit and neuronal differentiation 1 (CEND1) also known as BM88, is a neuronal protein associated in vivo with terminal neuron-generating divisions, marking the exit of proliferative cells from the cell cycle (PMID: 17971443). CEND1 expression is critical for the NEUROG2-driven reprogramming of astrocytes, suggesting the existence of a reciprocal feedback loop leading to neurogenesis (PMID: 26321141). CEND1 is an integral membrane protein composed of two 23-kDa polypeptide chains linked together by disulfide bridges.

cell cycle exit and neuronal

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 13280-1-AP (CEND1 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed Neuro-2a cells using CEND1 antibody (13280-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 13280-1-AP (CEND1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).