

For Research Use Only

RNF125 Polyclonal antibody

Catalog Number: 13290-1-AP

3 Publications



Basic Information

Catalog Number:

13290-1-AP

Size:

150ul, Concentration: 800 ug/ml by Nanodrop and 520 ug/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG3908

GenBank Accession Number:

BC012021

GeneID (NCBI):

54941

UNIPROT ID:

Q96EQ8

Full Name:

ring finger protein 125

Calculated MW:

232 aa, 26 kDa

Observed MW:

22-26 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IHC 1:20-1:200

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IHC

Species Specificity:

human

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : A2780 cells,

IHC : human lymphoma tissue,

Notable Publications

Author	Pubmed ID	Journal	Application
Xianghui Kong	36662617	Cell Rep	WB
Xueyi Feng	38298426	Oncol Lett	WB
Zhigang Feng	37142680	Oncogene	WB,IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



A2780 cells were subjected to SDS PAGE followed by western blot with 13290-1-AP (RNF125 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human lymphoma tissue slide using 13290-1-AP (RNF125 Antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human lymphoma tissue slide using 13290-1-AP (RNF125 Antibody) at dilution of 1:50 (under 40x lens).