

For Research Use Only

ICOS Polyclonal antibody

Catalog Number: 13338-1-AP



Basic Information

Catalog Number: 13338-1-AP	GenBank Accession Number: BC028210	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 700 µg/ml by Nanodrop and 367 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 29851	Recommended Dilutions: WB 1:200-1:1000
Source: Rabbit	Full Name: inducible T-cell co-stimulator	
Isotype: IgG	Calculated MW: 199 aa, 23 kDa	
Immunogen Catalog Number: AG4155	Observed MW: 28 kDa, 45-50 kDa	

Applications

Tested Applications: WB,ELISA	Positive Controls: WB : Jurkat cells, HL-60 cells
Species Specificity: human	

Background Information

Inducible T cell co-stimulator (ICOS) is related to the CD28 superfamily and is highly expressed on activated T cells as well as regulatory T cells and is crucial for the survival and function of T cells, Th2 cell differentiation and for lung inflammatory responses. Binding ICOS to ICOS-ligand (ICOS-L) activates a cascade of intracellular signaling molecules that prevent apoptosis and lead to the production of cytokines such as IL-4 and IL-13. Catalog# 13338-1-AP recognises two band of 25-30 kDa and 45-50 kDa, and the additional 45-50 kDa band may be due to surfacedisulfide-linked homodimeric glycoprotein.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with 13338-1-AP (ICOS Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.