

For Research Use Only

NCAPD2 Polyclonal antibody

Catalog Number: 13382-1-AP

Featured Product

3 Publications



Basic Information

Catalog Number:
13382-1-AP

Size:
150ul, Concentration: 350 µg/ml by Nanodrop and 200 µg/ml by Bradford method using BSA as the standard;

Source:
Rabbit

Isotype:
IgG

Immunogen Catalog Number:
AG4219

GenBank Accession Number:
BC028182

GeneID (NCBI):
9918

Full Name:
non-SMC condensin I complex, subunit D2

Calculated MW:
1401 aa, 157 kDa

Observed MW:
157 kDa

Purification Method:
Antigen affinity purification

Recommended Dilutions:
WB 1:500-1:3000
IP 0.5-4.0 ug for IP and 1:500-1:1000 for WB
IF 1:20-1:200

Applications

Tested Applications:
IF, IP, WB, ELISA

Cited Applications:
IHC, WB

Species Specificity:
human, mouse, rat

Cited Species:
human, mouse

Positive Controls:

WB: HEK-293 cells, COLO 320 cells, HeLa cells, K-562 cells

IP: HEK-293 cells,

IF: HeLa cells,

Background Information

NCAPD2 is a regulatory subunit of the condensin complex, which is required for conversion of interphase chromatin into mitotic-like condensed chromosomes. Condensin complexes are heteropentamers comprised of two SMC (structural maintenance of chromosomes) subunits and three non-SMC subunits. The C-terminus of NCAPD2 interacts with Histones H1 and H3 through their histone tails. A loss of NCAPD2 can lead to the disorganization of chromatid axes, misalignment of sister chromatids during metaphase and delayed entry into anaphase.

Notable Publications

Author	Pubmed ID	Journal	Application
Biao Huang	31704649	iScience	WB
Zuolei Jing	34229059	Cancer Lett	WB, IHC
Xiaoying Dong	37498296	Aging (Albany NY)	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



HEK-293 cells were subjected to SDS PAGE followed by western blot with 13382-1-AP (NCAPD2 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HeLa cells, using NCAPD2 antibody 13382-1-AP at 1:50 dilution and Rhodamine-labeled goat anti-rabbit IgG (red). Blue pseudocolor = DAPI (fluorescent DNA dye).



IP Result of anti-NCAPD2 (IP:13382-1-AP, 4ug; Detection:13382-1-AP 1:600) with HEK-293 cells lysate 2800ug.