For Research Use Only

KIF22 Polyclonal antibody

Catalog Number: 13403-1-AP

Featured Product

3 Publications

BC028155



Basic Information

Catalog Number: GenBank Accession Number:

GeneID (NCBI):

150ul, Concentration: 400 µg/ml by 3835 Nanodrop;

Source: kinesin family member 22

Rabbit Calculated MW: Isotype: 665 aa, 73 kDa IgG Observed MW: Immunogen Catalog Number: 68-73 kDa

13403-1-AP

AG4254

Antigen affinity purification Recommended Dilutions: WB 1:500-1:2000

Purification Method:

IP 0.5-4.0 ug for IP and 1:200-1:1000

for WB IHC 1:20-1:200

Applications

Tested Applications:

IHC, IP, WB, ELISA

Cited Applications: IF, IHC, WB

Species Specificity:

human, mouse, rat **Cited Species:**

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: Jurkat cells, human heart tissue, MKN-45 cells, T-

47D cells

IP: HeLa cells,

IHC: human cervical cancer tissue, human breast

cancer tissue

Background Information

KIF22, also known as KID (kinesin-like DNA-binding protein), is a kinesin-like DNA-binding protein known to be involved in chromosome movement during mitosis. KIF22 is a nuclear protein regulated by SIAH-1, whose level fluctuate in a cell cycle-dependent manner, increasing during pre-mitotic phases and greatly decreasing during mitosis. KIF22 localizes to the nucleus of the interphase cells upon entry into mitosis and displays some punctate cytoplasmic staining. Defects in KIF22 are the cause of spondyloepimetaphyseal dysplasia with joint laxity, type 2 (SEMDJL2).

Notable Publications

Author	Pubmed ID	Journal	Application
Sara Bizzotto	29229923	Sci Rep	IF
Yan Guang-Rong GR	22887948	Proteomics	WB
Ze-Yuan Yu	32187050	Chin Med J (Engl)	WB,IHC

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 13403-1-AP (KIF22 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human cervical cancer using 13403-1-AP (KIF22 antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human cervical cancer using 13403-1-AP (KIF22 antibody) at dilution of 1:100 (under 40x lens)



IP Result of anti-KIF22 (IP:13403-1-AP, 4ug; Detection:13403-1-AP 1:300) with HeLa cells lysate 4000ug.