

For Research Use Only

VASP Polyclonal antibody

Catalog Number: 13472-1-AP

Featured Product

9 Publications



Basic Information

Catalog Number: 13472-1-AP	GenBank Accession Number: BC038224	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 500 µg/ml by Nanodrop;	GeneID (NCBI): 7408	Recommended Dilutions: WB 1:1000-1:6000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:20-1:200 IF 1:50-1:500
Source: Rabbit	Full Name: vasodilator-stimulated phosphoprotein	
Isotype: IgG	Calculated MW: 380 aa, 40 kDa	
Immunogen Catalog Number: AG4266	Observed MW: 46 kDa, 50 kDa	

Applications

Tested Applications:
FC, IF, IHC, IP, WB, ELISA

Cited Applications:
IF, IHC, WB

Species Specificity:
human, mouse, rat

Cited Species:
human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A549 cells, mouse spleen tissue, HEK-293 cells, HeLa cells, C6 cells

IP: HEK-293 cells,

IHC: human testis tissue, human heart tissue, human kidney tissue, human lung tissue, human skin cancer tissue, human spleen tissue

IF: HepG2 cells,

Background Information

VASP belongs to the Ena/VASP family. Ena/VASP proteins are actin-associated proteins involved in a range of processes dependent on cytoskeleton remodeling and cell polarity such as axon guidance, lamellipodial and filopodial dynamics, platelet activation and cell migration. VASP promotes actin filament elongation. It protects the barbed end of growing actin filaments against capping and increases the rate of actin polymerization in the presence of capping protein. VASP stimulates actin filament elongation by promoting the transfer of profilin-bound actin monomers onto the barbed end of growing actin filaments. VASP plays a role in actin-based mobility of *Listeria monocytogenes* in host cells. Regulates actin dynamics in platelets and plays an important role in regulating platelet aggregation. Human platelet activation is inhibited by agents such as prostaglandins and NO donors, which elevate cAMP or cGMP levels. VASP is phosphorylated in human platelets in response to both cAMP- and cGMP-elevating agents, and its phosphorylation correlates with platelet inhibition. VASP is located about 92 kb distal to ERCC1 (126380) and about 300 kb proximal to the myotonic dystrophy protein kinase gene. The antibody is specific to VASP.

Notable Publications

Author	Pubmed ID	Journal	Application
Thibault Courtheoux	27641145	Nat Commun	IF
Jun Wang	34745942	Front Oncol	WB
Yajing Lv	33186350	PLoS Biol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 13472-1-AP (VASP antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human testis using 13472-1-AP (VASP antibody) at dilution of 1:100 (under 10x lens).




Immunohistochemical analysis of paraffin-embedded human testis using 13472-1-AP (VASP antibody) at dilution of 1:100 (under 40x lens).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using VASP antibody (13472-1-AP) at dilution of 1:200 and CoralLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IP result of anti-VASP(IP:13472-1-AP, 4ug; Detection:13472-1-AP 1:10000) with HEK-293 cells lysate 920 ug.



1X10⁶ HepG2 cells were stained with 0.2ug VASP antibody (13472-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.