For Research Use Only

p115, USO1 Polyclonal antibody

Catalog Number: 13509-1-AP

Featured Product

25 Publications



Basic Information

Catalog Number: GenBank Accession Number:

13509-1-AP BC032654 GeneID (NCBI): Size:

150ul , Concentration: 550 ug/ml by

Nanodrop: **UNIPROT ID:** 060763 Rabbit Full Name:

Isotype: USO1 homolog, vesicle docking

protein (yeast) IgG Immunogen Catalog Number: Calculated MW: 962 aa, 108 kDa AG4431

> Observed MW: 115 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IF, CoIP Species Specificity: human, mouse, rat **Cited Species:**

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:8000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF/ICC 1:500-1:2000

human, mouse

Positive Controls:

WB: HEK-293 cells, mouse testis tissue, mouse thymus tissue, HeLa cells, human brain tissue, SH-SY5Y cells,

HepG2 cells

IP: mouse brain tissue, IHC: human gliomas tissue,

IF/ICC: HeLa cells,

Background Information

p115, also known as USO1, TAP (transcytosis-associated protein) or VDP (vesicle docking protein) is a general vesicular transport factor and plays an important role at different steps of vesicular transport. It is a 962-residue peripheral membrane protein which recycles between the cytosol and the Golgi apparatus during interphase (PMID: 9478999). p115 forms stable homodimers (PMID: 19247479). Rab1 recruits p115 to coat protein complex II (COPII) vesicles during budding from the endoplasmic reticulum, where p115 interacts directly with a select set of SNARE proteins (PMID: 10903204). p115 is required for intra-Golgi transport, and also functions in endoplasmic reticulum to Golgi trafficking, Golgi biogenesis and exocytotic transport (PMID: 19247479).

Notable Publications

Author	Pubmed ID	Journal	Application
Mohsan Saeed	32997711	PLoS Pathog	WB
Guillermo Arango Duque	34580108	J Immunol	IF
Jing Wang	29025970	J Cell Sci	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



WB result of p115, USO1 antibody (13509-1-AP; 1:40000; incubated at room temperature for 1.5 hours) with sh-Control and sh-p115, USO1 transfected HeLa cells.




Various lysates were subjected to SDS PAGE followed by western blot with 13509-1-AP (p115, USO1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed HeLa cells using p115, USO 1 antibody (13509-1-AP) at dilution of 1:1000 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 13509-1-AP (p115, USO 1 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 13509-1-AP (p115, USO1 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-p115, USO1 (IP:13509-1-AP, 3ug; Detection:13509-1-AP 1:1500) with mouse brain tissue lysate 7000ug.