## For Research Use Only

## BST2 Polyclonal antibody, PBS Only

Catalog Number:13560-1-PBS Featured Product



**Purification Method:** 

Antigen affinity purification

**Basic Information** 

Catalog Number: GenBank Accession Number:

13560-1-PBS BC03387

Size: GeneID (NCBI):

100ug, Concentration: 1 mg/ml by 684

Nanodrop; UNIPROT ID:
Source: Q10589
Rabbit Full Name:

Isotype: bone marrow stromal cell antigen 2

lgG Calculated MW:
Immunogen Catalog Number: 180 aa, 20 kDa
AG4430 Observed MW:
30-36 kDa, 50-70 kDa

Applications Tested Applications:

WB, IHC, IF/ICC, Indirect ELISA

Species Specificity:

human


## **Background Information**

BST2, also named as CD317 and Tetherin, belongs to the tetherin family. It may be involved in the sorting of secreted proteins and it is involved in pre-B-cell growth. BST2 is an antiretroviral defense protein, that blocks retrovirus release from the cell surface. Depleted upon HIV-1 infection by viral VPU protein through 20S proteasome degradation. Depleted upon infection by human Kaposi's sarcoma-associated herpesvirus (KSHV) through ubiquitination and subsequent degradation. BST2 may play a role in B-cell activation in rheumatoid arthritis. It is recently identified interferon-induced cellular proteins that restrict infections by retroviruses and filoviruses and of influenza virus and flaviviruses, respectively. BST2 is a plasma membrane protein, tetherin inhibits virion particle release from infected cells. BST2 is effective against retroviruses and flavoviruses whilst IFITMs disrupt influenza and flavivirus infection. Observed MW of BST2 are 30-36 kDa (monomer form) and 50-70 kDa (dimer form) (PMID: 19196977; 21237475; 19737401; 37922253).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 13560-1-AP (BST2 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 13560-1-PBS in a different storage buffer formulation.



WB result of BST2 antibody (13560-1-AP; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-BST2 transfected HeLa cells. This data was developed using the same antibody clone with 13560-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 13560-1-AP (BST2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 13560-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed NCCIT cells using BST2 antibody (13560-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 13560-1-PBS in a different storage buffer formulation.