

For Research Use Only

PRDX6 Polyclonal antibody

Catalog Number: 13585-1-AP

Featured Product

35 Publications



Basic Information

Catalog Number:

13585-1-AP

Size:

150ul, Concentration: 550 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG4512

GenBank Accession Number:

BC035857

GeneID (NCBI):

9588

UNIPROT ID:

P30041

Full Name:

peroxiredoxin 6

Calculated MW:

224 aa, 25 kDa

Observed MW:

25-30 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:12000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:20-1:200

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: SH-SY5Y cells, A549 cells, HeLa cells, HAP1, K-562 cells, HepG2 cells, mouse liver tissue, rat liver tissue

IP: HeLa cells, HAP1 cells

IHC: human brain tissue,

IF/ICC: 48 HepG2 cells, HeLa cells

Background Information

PRDX6 (Peroxiredoxin-6), also named as AOP2 or KIAA0106, is a unique member of the peroxiredoxin family of antioxidants. PRDX6 is highly expressed in liver and protects cells from oxidative damage by reducing H₂O₂ and various lipid Peroxides (PMID: 17382207). It can form a dimer (PMID: 20500660). PRDX6 is expressed in all major organs, with a particularly high level in lung (PMID: 15890616). Prdx6 is detected at approximately 24 to 28 kDa, and can be monosumoylated with the molecular mass of about 40 kDa (PMID: 24910119).

Notable Publications

Author	Pubmed ID	Journal	Application
Jiale Sun	34557413	Front Oncol	IHC
Chenhao Xu	33161253	Bioorg Chem	WB
Siddhesh Aras	25315652	Mitochondrion	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 13585-1-AP (PRDX6 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




WB result of PRDX6 antibody (13585-1-AP; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PRDX6 transfected HeLa cells.




HAP1 (WT and PRDX6 KO) lysates prepared with RIPA buffer, 25 µg protein loaded. 13585-1-AP incubated at 1:2000 at 4°C overnight in 5% BSA in TBST. Ponceau stained transfers shown on right. Data provided by YCharOS, an open science company with a mission to validate commercial antibodies to improve scientific reproducibility and transparency.




Various lysates were subjected to SDS PAGE followed by western blot with 13585-1-AP (PRDX6 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




IP result of anti-PRDX6 (IP:13585-1-AP, 3ug; Detection:13585-1-AP 1:500) with HeLa cells lysate 2800ug.




HAP1 lysates prepared and IP of peroxidoxin 6 performed using 2.0 µg of 13585-1-AP coupled to protein A- Sepharose beads. Ponceau stained transfers shown for each blot. Data provided by YCharOS, an open science company with a mission to validate commercial antibodies to improve scientific reproducibility and transparency.






HAP1 WT cells (yellow outline) and PRDX6 KO cells (red outline) labelled with a green or a far red fluorescence dye, respectively. Cells fixed with 4% PFA and stained with 13585-1-AP at 1:300 plus DAPI. Bars = 10 µm. Data provided by YCharOS, an open science company with a mission to validate commercial antibodies to improve scientific reproducibility and transparency.




Immunohistochemical analysis of paraffin-embedded human brain using 13585-1-AP (PRDX6 antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human brain using 13585-1-AP (PRDX6 antibody) at dilution of 1:50 (under 40x lens).



Immunohistochemical analysis of paraffin-embedded human normal colon slide using 13585-1-AP (PRDX6 antibody) at dilution of 1:1200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed 48 HepG2 cells using PRDX6 antibody (13585-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).

Immunohistochemical analysis of paraffin-embedded human normal colon slide using 13585-1-AP (PRDX6 antibody) at dilution of 1:1200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using PRDX6 antibody (13585-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).