For Research Use Only

## SUR2/ABCC9 Polyclonal antibody

Catalog Number:13599-1-AP




Basic Information	Catalog Number: 13599-1-AP	GenBank Accession Number: BC033804	Purification Method: Antigen affinity purification
	Size: 150ul, Concentration: 650 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG4536	GeneID (NCBI): 10060 UNIPROT ID: 060706 Full Name: ATP-binding cassette, sub-family C	Recommended Dilutions: WB 1:500-1:1000
		Calculated MW: 1549 aa, 174 kDa	
		Observed MW: 174 kDa	
		Applications	Tested Applications: WB, ELISA
Species Specificity: human, mouse			
Background Information	ABCC9, also known as SUR2, belongs to the ABC transporter superfamily. ABCC9 is a subunit of ATP-sensitive potassium channels (KATP), forming cardiac and smooth muscle-type KATP channels with KCNJ11. KCNJ11 forms the channel pore while ABCC9 is required for activation and regulation (PMID: 9831708). ABCC9 is thought to form ATP-sensitive potassium channels in cardiac, skeletal, vascular, and non-vascular smooth muscle. Defects in ABCC9 cause dilated cardiomyopathy 10 and familial atrial fibrillation type 12 (ATFB12)(PMID: 15034580,17245405).		
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3.		
		torage	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free<br/>in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 13599-1-AP (SUR2/ABCC9 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



mouse heart were subjected to SDS PAGE followed by western blot with 13599-1-AP (SUR2/ABCC9 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.